

(12) **EUROPEAN PATENT APPLICATION**


(21) Application number: **84201851.7**
 (22) Date of filing: **13.12.84**

(51) Int. Cl.⁴: **C 07 D 471/04**
C 07 D 401/06, C 07 D 401/12
C 07 D 401/14, C 07 D 405/14
C 07 D 409/14, C 07 D 417/14
C 07 D 513/04, C 07 D 519/00
C 07 D 487/04, C 07 D 473/00

(30) Priority: **09.01.84 US 569369**
13.11.84 US 571135
 (43) Date of publication of application:
21.08.85 Bulletin 85/34
 (84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(71) Applicant: **JANSSEN PHARMACEUTICA N.V.**
Turnhoutsebaan 30
B-2340 Beerse(BE)
 (72) Inventor: **Janssens, Frans Eduard**
Tinstraat 79
B-2830 Bonheiden(BE)
 (72) Inventor: **Kennis, Ludo Edmond Josephine**
Guido Gezellenstraat 50
B-2300 Turnhout(BE)
 (72) Inventor: **Hens, Jozef Francis**
Guido Gezellestraat 50
B-2260 Nijlen(BE)
 (72) Inventor: **Torremans, Joseph Leo G.**
Lijsterstraat 11
B-2340 Beerse(BE)
 (72) Inventor: **Diels, Gaston Stanislas M.**
Oosteinde 12
B-2380 Ravels(BE)

(54) **4-(Bicyclic heterocyclyl)-methyl and -hetero-piperidines.**
 (57) Novel 4-[(bicyclic heterocyclyl)methyl and -hetero]-piperidines of formula



wherein B is CH₂, O, S, SO or SO₂; the pharmaceutically acceptable acid addition salts and the possible stereochemically isomeric forms thereof, which compounds are anti-allergic agents, pharmaceutical compositions containing such compounds as an active ingredient and processes for preparing the said compounds and compositions

EP 0 151 826 A1

-1-

5

10


4-[(BICYCLIC HETEROCYCLYL)-METHYL AND -HETERO]-PIPERIDINES.

15

Background of the invention:

In U.S. Patent No. 4,219,559 there are described a number of
N-heterocyclyl-4-piperidinamines having the formula

20




25

which compounds are useful as antihistaminic agents.

The compounds of the present invention differ from the prior art compounds essentially by the nature of the 4-piperidinyl substituent which is invariably a bicyclic heterocyclyl-methyl or -hetero group and by the fact that the compounds of the present invention are not
30 only potent histamine-antagonists but also potent serotonin-antagonists.

Description of the preferred embodiments:

This invention is concerned with novel 4-[bicyclic heterocyclyl-methyl and -hetero]-piperidines which may structurally be represented
35 by the formula



5

the pharmaceutically acceptable acid addition salts and the possible stereochemically isomeric forms thereof, wherein:

$A^1-A^2-A^3-A^4$ is a bivalent radical having the formula

- 10 $-CH=CH-CH=CH-$ (a-1),
 $-N=CH-CH=CH-$ (a-2),
 $-CH=N-CH=CH-$ (a-3),
 $-CH=CH-N=CH-$ (a-4), or
 $-CH=CH-CH=N-$ (a-5),

15 wherein one or two hydrogen atoms in said radicals (a-1) - (a-5) may, each independently from each other, be replaced by halo, lower alkyl, lower alkyloxy, trifluoromethyl or hydroxy;

R^1 is a member selected from the group consisting of hydrogen, alkyl, cycloalkyl, Ar^1 and lower alkyl substituted with one or two Ar^1 radicals;


20 R^2 is a member selected from the group consisting of hydrogen and lower alkyl;

 B is CH_2 , O, S, SO or SO_2 ;

 L is a member selected from the group consisting of a radical of formula



a radical of formula



30

wherein one or two hydrogen atoms in the bivalent radical

$-C_{sH_{2s}}-$ may, each independently from each other, be replaced by halo, hydroxy, mercapto, isothiocyanato, isocyanato, lower alkyloxy, lower alkylthio, Ar^1 , Ar^1O- , Ar^1S- , Ar^1SO_2- , or NR^3R^5 ; and

35 n is 0 or the integer 1 or 2;

r and s are, independently from each other, 0 or an integer of from 1 to 6 inclusive;

T is -Y- or $-Z-\overset{\overset{X}{\parallel}}{C}-Y-$;

5

T¹ is $-\overset{\overset{X}{\parallel}}{Z}-C-$ or a direct bond;

said Y being O, S, NR³ or a direct bond;

10 X being O, S, CH-NO₂ or NR⁴;

Z being O, S, NR⁵ or a direct bond; and

said R³ being hydrogen, lower alkyl, (Ar²)lower alkyl, 2-lower alkyloxy-1,2-dioxoethyl or a radical of formula -C(=X)-R⁶, R⁶ being hydrogen, lower alkyl, Ar², Ar²-lower alkyl, lower alkyloxy,

15 Ar²-lower alkyloxy, mono- or di(lower alkyl)amino, Ar²-amino, Ar²-lower alkylamino or Ar²-lower alkyl(lower alkyl)amino;

said R⁴ being hydrogen, lower alkyl, cyano, nitro, Ar²-sulfonyl, lower alkylsulfonyl, lower alkylcarbonyl or Ar²-carbonyl; and


said R⁵ being hydrogen or lower alkyl;

20

wherein L¹ is a member selected from the group consisting of hydrogen; halo; hydroxy; lower alkyloxy; lower alkylthio; cyano; mercapto; isocyanato; isothiocyanato; Ar¹; Ar¹-carbonyl; Ar¹-sulfonyl; lower alkylsulfonyl; cycloalkyl being optionally substituted with up to

25 two substituents each independently selected from the group consisting of lower alkyl, cyano and Ar²; [10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene]methyl; Het; and furan substituted with substituted lower alkyl; said substituted lower alkyl being lower alkyl substituted with a member selected from the group consisting of hydroxy, mercapto,

30 lower alkyloxy, lower alkylthio, aminolower alkylthio, Ar²-oxy and a radical of formula



(c),

wherein: t is 0 or an integer of from 1 to 6 inclusive; and
 R^7 is hydrogen or lower alkyl;
 provided that: when in said radical of formula (c) t is
 0, then Z or Y is a direct bond; and

5

where r is 0, L^1 may also be lower alkenyl, Ar^1 -lower alkenyl or
 lower alkyl substituted with two lower alkyloxy radicals; and

where r is 0 and T is NR^3 , or T is $-N(R^5)-C(=X)-Y$ or T^1 is $-N(R^5)-C(=X)-$,
 10 L^1 may also be amino, lower alkylamino or Ar^1 -amino; and

where r is 0, and T is $-N(R^5)-C(=X)-Y$ or T^1 is $-N(R^5)-C(=X)-$,
 L^1 may also be nitro;

15 said Het being an optionally substituted five- or six-membered hetero-
 cyclic ring, being optionally condensed with an optionally substituted
 five- or six-membered carbocyclic or heterocyclic ring;

provided that:

- 20 i) when L is a radical of formula (b-1) wherein L^1 is hydrogen and
 wherein T is $-Z-C(=X)-Y-$ wherein Y is other than a direct bond
 and Z and X are each independently O or S, then r is not 0; or
 when L is a radical of formula (b-2) wherein L^1 is hydrogen and
 wherein T^1 is $-Z-C(=X)-$ wherein Z and X are each independently
 25 O or S, then r is not 0;
- ii) when L is a radical of formula (b-1) wherein L^1 is halo,
 hydroxy, lower alkyloxy, mercapto, lower alkylthio, isocyanato,
 isothiocyanato or Het connected to $C_{r-2r}H_{2r}$ on a nitrogen atom,
 and wherein r is 0, then T is a direct bond or a radical
 30 $-C(=X)-Y-$; or when L is a radical of formula (b-2) wherein L^1
 is halo, hydroxy, lower alkyloxy, mercapto, lower alkylthio,
 isocyanato, isothiocyanato or Het connected to $C_{r-2r}H_{2r}$ on a
 nitrogen atom, and wherein r is 0, then T^1 is a radical $-C(=X)-$;

35

iii) when L is a radical of formula (b-1) wherein T is Y, said Y being other than a direct bond, or wherein T is $-Z-C(=X)-Y-$, wherein Y is other than a direct bond, then s is not 0;

- 5 wherein Ar^1 is a member selected from the group consisting of phenyl, substituted phenyl, naphthalenyl, thienyl, halothienyl, lower alkylthienyl, pyridinyl, mono- and di(lower alkyloxy)pyridinyl, pyrrolyl, lower alkylpyrrolyl, furanyl, furanyl substituted with lower alkyl, pyrazinyl, thiazolyl, imidazolyl, lower alkylimidazolyl;
- 10 said substituted phenyl, being phenyl substituted with up to 3 substituents each independently selected from the group consisting of halo, hydroxy, nitro, cyano, trifluoromethyl, lower alkyl, lower alkyloxy, lower alkylthio, mercapto, amino, mono- and di(lower alkyl)amino, lower alkylsulfonyl, lower alkylsulfonylower alkyl,
- 15 phenyllower alkylsulfonyl, phenylsulfonylower alkyl, a radical of formula $R^8-C_{p2p}H_{2p}-Y-$, a radical of formula $R^9-Z-C(=X)-Y-$, and a radical of formula $R^{10}SO_2Y-$; wherein p is an integer of from 1 to 6 inclusive and R^8 is a member selected from the group consisting of amino, cyano, phenyl aminocarbonyl, mono- and di(lower alkyl)amino-
- 20 carbonyl, lower alkyloxycarbonyl, phenyllower alkyloxycarbonyl, 4-morpholinylcarbonyl, 1-piperidinylcarbonyl, 1-pyrrolidinylcarbonyl, and lower alkenyl; wherein R^9 is member selected from the group consisting of hydrogen, lower alkyl and Ar^2 ; provided that, when R^9 is hydrogen and Y is other than a direct bond, then Z is not O or
- 25 S; and wherein R^{10} is lower alkyl or Ar^2 ;

wherein Ar^2 is a member selected from the group consisting of phenyl, substituted phenyl, thienyl and furanyl, said substituted phenyl being phenyl optionally substituted with up to three substituents each independently selected from the group consisting of

30 halo, hydroxy, nitro, cyano, trifluoromethyl, lower alkyl, lower alkyloxy, lower alkylthio, mercapto, amino, mono- and di(lower alkyl)amino, carboxyl, lower alkyloxycarbonyl and (lower alkyl)-CO.

As used in the foregoing definitions the term hal is generic to fluoro, chlor, bromo and iodo; the term "lower alkyl" is meant to include straight and branch chained saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as, for example, methyl, ethyl, 5 1-methylethyl, 1,1-dimethylethyl, propyl, 2-methylpropyl, butyl, pentyl, hexyl and the like; "alkyl" is meant to include lower alkyl radicals, as defined hereinabove, and the higher homologs thereof having from 7 to 10 carbon atoms; the term "lower alkenyl" is meant to include straight and branch chained hydrocarbon radicals having from 2 10 to 6 carbon atoms, such as, for example, 2-propenyl, 2-butenyl, 3-butenyl, 2-pentenyl and the like; the term "cycloalkyl" is generic to cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; and "lower alkanediyl" is meant to include bivalent straight or branch chained alkanediyl radicals having from 1 to 6 carbon atoms.

15 Preferred compounds within the invention are those wherein r is 0 and L^1 is hydrogen, hydroxy, lower alkyloxy, lower alkylthio, mercapto, Het, Ar^1 , cyanato, isocyanato or isothiocyano.

Particularly preferred compounds within the invention are those wherein r is 0 and L^1 is as described hereinabove for the preferred 20 compounds and wherein R^1 is lower alkyl substituted with one Ar^1 radical.


More particularly preferred compounds within the invention are those wherein L is a radical of formula (b-1), wherein r is 0 and L^1 is as described hereinabove for the preferred compounds and wherein 25 R^1 is lower alkyl substituted with one Ar^1 radical.

It is evident that in the compounds of formula (I) wherein L^1 is Het, said Het may be unsaturated or partly or completely saturated.

The compounds of formula (I) wherein Het is a heterocycle which is substituted with a hydroxy, mercapto or amino radical may contain in 30 their structure a keto-enol tautomeric system or a vinylog system thereof, and consequently these compounds may be present in their keto form as well as their enol form.

The compounds of formula (I) can generally be prepared by reacting

a piperidine of formula (II) with a diamine of formula (III).



In (II) X^1 is O, S or NH.

W as used in the foregoing and following reaction schemes is an appropriate leaving group such as, for example, halo, e.g. chloro, bromo or iodo, a sulfonyloxy group, e.g. methylsulfonyloxy or 4-methylphenylsulfonyloxy, and where W is connected to a $-C(=X)-$, $-C(=X^1)-$ or $-C(=X^2)-$ radical it may also be lower alkyloxy, lower alkylthio, Ar^2-O- , or Ar^2-S- .

15 The piperidine of formula (II) may in situ be generated, for example, by converting a piperidine which is substituted in its 4-position with a $-B-C(=X^1)-OH$ radical into a piperidine of formula (II) by reacting the former piperidine with thionyl chloride, phosphor trichloride, polyphosphoric acid, phosphoroxychloride and the like.


20 The reaction of (II) with (III) may be conducted in a suitable solvent such as, for example, a hydrocarbon, e.g., benzene, hexane, an ether, e.g., 1,1'-oxybisethane, tetrahydrofuran, a ketone, e.g., propanone, an alcohol, e.g., methanol, ethanol, 2-propanol, 1-butanol, a halogenated hydrocarbon, e.g., trichloromethane, dichloromethane, an

25 acid, e.g., acetic acid, propanoic acid, N,N-dimethylformamide, N,N-dimethylacetamide and the like, and mixtures of such solvents.


Depending upon the solvent and nature of W it may be appropriate to add an appropriate base and/or an iodide salt, preferably an alkali metal iodide, to the reaction mixture. Elevated temperatures may

30 enhance the reaction rate.


The compounds of formula (I) can also be prepared by reacting an intermediate of formula (V) with a piperidine of formula (IV) wherein E^1 and E^2 are selected so that during the reaction a radical $-B-$ is formed.



For example, the compounds of formula (I) can be prepared by reacting a piperidine of formula (IV) wherein E^1 is a radical of formula $-B-M$ with an intermediate of formula (V) wherein E^2 is a radical of formula $-W$.




In (IV-a) M is, depending upon the nature of B , hydrogen or an appropriate alkali metal or earth alkaline metal and in (V-a) W has the previously described meaning. Additionally, the compounds of formula (I) can also be prepared by reacting a piperidine of formula (IV) wherein E^1 is W with an intermediate of formula (V) wherein E^2 is a radical of formula $-B-M$, said W and M having the previously described meanings.



More particularly, the compounds of formula (I) wherein B is $-\text{CH}_2-$ can also be prepared by reacting a piperidine of formula (IV) wherein E^1 represents a radical of formula $-\text{CH}_2-W$, (IV-c), with an

intermediate of formula (V) wherein E^2 represents M, (V-c) or alternatively, by reacting a piperidine of formula IV, wherein E^1 is a radical of formula -M, (IV-d), with an intermediate of formula (V) wherein E^2 is a radical of formula $-CH_2-W$, (V-d).

5



20 The reaction of (IV) with (V) may conveniently be conducted in an appropriate solvent such as for example, an aromatic hydrocarbon, e.g., benzene, methylbenzene; an ether, e.g. 1,4-dioxane, 1,1'-oxybisethane, tetrahydrofuran and the like; a halogenated hydrocarbon, e.g. trichloromethane and the like; N,N-dimethylformamide (DMF); N,N-dimethylacet-


25 amide (DMA); and where M is hydrogen, said solvent may also be a lower alkanol, e.g., methanol, ethanol, 1-butanol and the like; a ketone, e.g., 2-propanone, 4-methyl-2-pentanone and the like. In some circumstances, the addition of an appropriate base such as, for example, an alkali metal carbonate or hydrogen carbonate, sodium hydride or an

30 organic base such as, for example, N,N-diethylethanamine or N-(1-methylethyl)-2-propanamine and/or the addition of an iodide salt, preferably an alkali metal iodide, may be appropriate. Somewhat elevated temperatures may enhance the rate of the reaction.

35

The compounds of formula (I) can also be derived from a 1,4-dihydropyridine derivative of formula (VI) following art-known reducing procedures.


5



10 Suitable reducing procedures are, for example, a catalytic hydrogenation in a suitable solvent, e.g. methanol, ethanol and the like, in the presence of a suitable catalyst, e.g. platinum-on-charcoal, palladium-on-charcoal and the like catalysts.

15 The compounds of formula (I) can also be converted into each other. A number of such reactions will be described hereinafter in more detail.

In order to simplify the structural representations of the compounds of formula (I) and of certain precursors and intermediates thereof the
20



25 represented by the symbol D.

The compounds of formula (I) wherein L is L², said compounds being represented by the formula (I-b) can be prepared by alkylating an intermediate of formula (VII) with a compound of formula (I) wherein L is Q², said compound being represented by the formula (I-c).




L^2 as defined hereinabove is a radical of formula (b-1) other than hydrogen, said radical being represented by the formula (b-1-a), or a radical of formula (b-2).




5

In (VII) and (I-c), Q^1 and Q^2 are selected so that a bivalent radical of formula (b-1-a) or (b-2) is formed during the alkylation reaction, said (b-1-a) and (b-2) having the previously described meaning.


10 For example, the compounds of formula (I-b) can be prepared by N-alkylating a piperidine of formula (I-c) wherein Q^2 is hydrogen, said piperidine being represented by the formula (I-c-1), with a reagent of formula (VII-a)



Additionally, the compounds of formula (I-b), wherein L^2 is a radical of formula (b-1-a), wherein T is T^2 , said T^2 being O, S, NR^3 or $-Z^1-C(=X)-Y-$, said Z^1 being O, S or NR^5 , or a radical of formula (b-2) wherein T^1 is T^3 , said T^3 being $-Z^1-C(=X)-$ or a direct bond, said compounds being represented by the formulae (I-b-1-a), respectively (I-b-1-b), can be prepared by alkylating a piperidine of formula (I-c-2) with a reagent of formula (VII-b).




35 In (I-c-2) Q^{2a} is a radical of formula $HT^2-C_{sH_{2s}}-$,

respectively a radical of formula HT^3-N 

and W^1 has the previously defined meaning of W , and where $r = 0$, and
 5 L^1 is Het or Ar^1 , it may also be lower alkyloxy or lower alkylthio.


The compounds of formula (I-b), wherein L^2 is a radical of
 formula (b-1-a), wherein T is T^4 , said T^4 being O, S, NR^3 or
 $-Z-C(=X)-Y^1-$, said Y^1 being O, S or NR^3 , and said compounds
 10 being represented by the formula (I-b-2), may also be prepared by
 alkylating a piperidine of formula (I-c) wherein Q^2 is a radical of
 formula $-C_{sH_{2s}}-W$, said piperidine being represented by the formula
 (I-c-3), with a reagent of formula (VII) wherein Q^1 is a radical of
 formula $-C_{rH_{2r}}-T^4H$, said reagent being represented by the
 15 formula (VII-c).




20

The alkylation reactions are conveniently conducted in an inert
 organic solvent such as, for example, an aromatic hydrocarbon, e.g.,
 benzene, methylbenzene, dimethylbenzene, and the like; a lower
 alkanol, e.g., methanol, ethanol, 1-butanol and the like;
 25 a ketone, e.g., 2-propanone, 4-methyl-2-pentanone and the like; an
 ether, e.g., 1,4-dioxane, 1,1'-oxybisethane, tetrahydrofuran and the
 like; N,N-dimethylformamide (DMF); N,N-dimethylacetamide (DMA);
 dimethyl sulfoxide (DMSO); nitrobenzene; 1-methyl-2-pyrrolidinone; and
 the like. The addition of an appropriate base such as, for example, an
 30 alkali metal carbonate or hydrogen carbonate, sodium hydride or an
 organic base such as, for example, N,N-diethylethanamine or
N-(1-methylethyl)-2-propanamine may be utilized to pick up the acid
 which is liberated during the course of the reaction. In some
 circumstances the addition of an iodide salt, preferably an alkali
 35 metal iodide, is appropriate. Somewhat elevated temperatures may

The compounds of formula (I-b) can also be prepared by the reductive N-alkylation reaction of (I-c-1) with an appropriate carbonyl-compound of formula $L^{2-a}C=O$ (VIII), said $L^{2-a}C=O$ being a compound of formula L^2-H wherein a $-CH_2-$ radical is oxidized to a carbonyl radical.



The compounds of formula (I-b), wherein L^2 is a radical of formula $L^1-C_{rH_{2r}}-NR^3-C_{sH_{2s}}-$, said compounds being represented by the formula (I-b-3) may alternatively be prepared by the reductive N-alkylation reaction of a compound of formula (I), wherein L is a radical of formula $HN(R^3)-C_{sH_{2s}}-$, (I-d), with an appropriate carbonyl-compound of formula $L^1-(C_{rH_{2r-1}})=O$, (IX), said $L^1-(C_{rH_{2r-1}})=O$ being a compound of formula $L^1-C_{rH_{2r}}-H$ wherein a $-CH_2-$ radical is oxidated to a carbonyl radical. The compounds of formula (I-b-3) can also be prepared by the reductive N-alkylation of an amine of formula (X), with a compound of formula (I) wherein L is a radical of formula $O=(C_{sH_{2s-1}})-$, said compound being represented by the formula (I-e), and said $O=(C_{sH_{2s-1}})-$ being a radical of formula $H-C_{sH_{2s}}-$ wherein a $-CH_2-$ radical is oxidated to a carbonyl radical.

$$\begin{array}{ccccc} \text{L}^1-(\text{C}_r\text{H}_{2r-1})=\text{O} + \text{HN}(\text{R}^3)-\text{C}_s\text{H}_{2s}-\text{D} & \longrightarrow & \text{L}^1-\text{C}_r\text{H}_{2r}-\text{N}(\text{R}^3)-\text{C}_s\text{H}_{2s}-\text{D} \\ \text{(IX)} & & \text{(I-d)} & & \text{(I-b-3)} \end{array}$$


BNSDOCID <EP 0151826A1 | >

solvents are, for example, water; low *r* alkanols, e.g. methanol, ethanol, 2-propanol and the like; cyclic ethers, e.g.

1,4-dioxane and the like; halogenated hydrocarbons, e.g.

trichloromethane and the like; *N,N*-dimethylformamide; dimethyl


5 sulfoxide and the like; or a mixture of 2 or more of such solvents.

The term "art-known catalytic hydrogenating procedures" means that the reaction is carried out under hydrogen atmosphere and in the presence of an appropriate catalyst such as, for example, palladium-on-charcoal, platinum-on-charcoal and the like. In order to prevent the

10 undesired further hydrogenation of certain functional groups in the reactants and the reaction products it may be advantageous to add an appropriate catalyst-poison to the reaction mixture, e.g., thiophene and the like.


The compounds of formula (I-b), wherein *L* is a radical of formula
15 (b-1-a) wherein *T* is $Z^1-C(=X^2)-NH-$, Z^1 being as previously described, X^2 being O or S, and said compounds being represented by the formula (I-b-4), can generally be prepared by reacting an isocyanate or isothiocyanate of formula (I-f) with a reagent of formula (XI):

20




25


The compounds of formula (I-b), wherein L^2 is a radical of formula (b-1-a), wherein *T* is $-NH-C(=X^2)-Y^1-$, Y^1 being as previously described, and the compounds of formula (I-b), wherein L^2 is a radical of formula (b-1-a), wherein *T* is $-NH-C(=X^2)-$ and *s* is
30 0, and the compounds of formula (I-b), wherein L^2 is a radical of formula (b-2), wherein T^1 is $-NH-C(=X^2)-$, said compounds being represented by the formula (I-b-5-a), respectively (I-b-5-b) and (I-b-5-c), can be prepared by reacting an isocyanate or isothiocyanate of formula (XII) with a piperidine of formula (I-c-4), respectively
35 (I-c-1) and (I-c-5).



5




10




15 The reaction of (XI) with (I-f) and of (XII) with (I-c-4), respectively (I-c-1) and (I-c-5) may be conducted in a suitable reaction-inert solvent such as, for example, a hydrocarbon, e.g., benzene, a ketone, e.g., acetone, a halogenated hydrocarbon, e.g., dichloromethane, trichloromethane, an ether, e.g., 1,1'-oxybisethane, 20 tetrahydrofuran and the like. Elevated temperatures may be suitable to enhance the rate of the reaction.


The compounds of formula (I-b), wherein L^2 is a radical of formula (b-1-a), wherein T is $-\text{C}(=\text{X}^2)-\text{Y}^1-$, and the compounds of 25 formula (I-b), wherein L is a radical of formula (b-1-a), wherein s is 0 and T is a radical of formula $-\text{C}(=\text{X}^2)-$, and the compounds of formula (I-b) wherein L^2 is a radical of formula (b-2), wherein T^1 is $-\text{C}(=\text{X}^2)-$, said compounds being represented by the formula (I-b-6-a), respectively (I-b-6-b) and (I-b-6-c), may be prepared by 30 reacting a piperidine of formula (I-c-4), respectively (I-c-1) and (I-c-5) with a reagent of formula (XIII).



35



5





10

The reaction of (XIII) with (I-c-4), respectively (I-c-1) and (I-c-5) may generally be conducted following art-known esterification- or amidation reaction-procedures. For example, the carboxylic acid may be converted into a reactive derivative, e.g., an anhydride or a carboxylic acid halide, which subsequently, is reacted with (I-c-4), (I-c-1) or (I-c-5); or by reacting (XIII) and (I-c-4), respectively (I-c-1) and (I-c-5) with a suitable reagent capable of forming amides or esters, e.g., dicyclohexylcarbodiimide, 2-chloro-1-methylpyridinium iodide and the like. Said reactions are most conveniently conducted in a suitable solvent such as, for example, an ether, e.g. tetrahydrofuran, a halogenated hydrocarbon, e.g. dichloromethane, trichloromethane or a polar aprotic solvent, e.g. N,N-dimethylformamide. The addition of a base, e.g. N,N-diethylethanamine may be appropriate.


The compounds of formula (I-b), wherein L^2 is a radical of formula (b-1-a) wherein T is $-Z^1-C(=X)-Y^1-$, and the compounds of formula (I-b), wherein L^2 is a radical of formula (b-1-a) wherein s is 0 and T is $-Z^1-(C=X)-$, and the compounds of formula (I-b), wherein L^2 is a radical of formula (b-2) wherein T^1 is $-Z^1-C(=X)-$, said compounds being represented by the formula (I-b-7-a), respectively (I-b-7-b) and (I-b-7-c), can also be prepared by reacting (XI) with (I-c-4), respectively (I-c-1) and (I-c-5) in the presence of an appropriate $>C=X$ generating agent.

35

-17-




5




10 An appropriate $\begin{array}{c} \diagup \\ C=X \\ \diagdown \end{array}$ generating agent is, for example, 1,1'-thiocarbonylbis[1H-imidazole], 1,1'-carbonylbis[1H-imidazole], carbonic dichloride, carbonothioic dichloride, urea, thiourea, trichloroacetyl chloride, and the like. The reaction of (XI) with (I-c-4), (I-c-1) or (I-c-5) is conveniently conducted in a suitable solvent, such as, for
15 example, a hydrocarbon, e.g., benzene, methylbenzene; an ether, e.g., 1,1'-oxybisethane, tetrahydrofuran; a halogenated hydrocarbon, e.g., dichloromethane, trichloromethane and the like. The addition of a base such as, for example, an alkali metal carbonate or hydrogen carbonate or an organic base, e.g., N,N-diethylethanamine and the like, may be
20 appropriate.

The compounds of formula (I-b) wherein L^2 is a radical of formula (b-1), wherein s is an integer of from 2 to 6 inclusive, said compounds being represented by the formula (I-g) can be prepared by reacting an appropriate alkene of formula (XIV) with a piperidine of
25 formula (I-c-1).



30 The compounds of formula (I-b) wherein L^2 is a radical of formula $L^1-C_rH_{2r}-T-C_{s'-2}H_{2s'-4}-CH(Y^1H)-CH_2-$, wherein s' is an integer of from 2 to 6 inclusive, said compounds being represented by the formula (I-h) may also be prepared by reacting a reagent of formula (XV) with a piperidine of formula (I-c-1).

35





5

The reactions of (XIV) with (I-c-1), and (XV) with (I-c-1) may be conducted by stirring and, if desired, heating the reactants together. The said reactions may be conducted in a suitable solvent such as, for example, an alkanone, e.g. 2-propanone, 4-methyl-2-propanone, an ether, e.g. tetrahydrofuran, 1,1'-oxybisethane, an alcohol, e.g. methanol, ethanol, 1-butanol, N,N-dimethylformamide, N,N-dimethylacetamide and the like.

It is evident that the radical "-lower alkenyl-", the corresponding "-lower alkanediyl-" radical and the radical $\text{C}_{2s'-2}\text{H}_{2s'-4}$ may bear the previously described substitutions of the radical $-\text{C}_s\text{H}_{2s}-$.


The compounds of formula (I) wherein L^1 is Het, said compounds being represented by the formula (I-i), may also be prepared following procedures for preparing ring systems which are known in the art or analogous procedures thereof. A number of such cyclization procedures will be described hereinafter.

The bivalent radical K used in the description of these cyclization reactions has the following meaning:




30 and the radicals (e-1), (e-2), (e-3), (e-4), (e-5), (e-6) and (e-7) also used in the description of these cyclization reactions have the following meaning :


35




(e-1) ;




(e-2) ;




(e-3),




(e-4),



(e-5),



(e-6), and




(e-7); wherein

15

X^2 has the previously defined meaning; and R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} and R^{20} are, each independently optional substituents of the said radicals (e-1) - (e-7) and where (e-1), (e-5) or (e-6) is connected to CH_2 by the nitrogen or carbon bearing R^{11} , R^{17} , R^{18} or R^{19} said R^{11} , R^{17} , R^{18} or R^{19} is absent; and G^1 , G^2 , G^3 , G^4 and G^5 , being each independently, optionally substituted bivalent radicals, are selected so that they form, combined with the five- or six-membered heterocycle to which they are attached, a bicyclic Het-system.

25 For example, the compounds of formula (I-i) wherein Het is an optionally substituted imidazolyl radical, said compounds being represented by the formula (I-i-1), can be prepared by the cyclization reaction of an appropriate N-(2,2-dilower alkyloxyethyl)imidamide derivative of formula (XVI).



30



wherein R^{21} , R^{22} and R^{23} are each independently optional substituents of the imidazole ring.


Said cyclization reaction may conveniently be conducted in a suitable solvent in the presence of an appropriate acid such as, for example, hydrochloric, hydrobromic and the like acids. Elevated temperatures may enhance the rate of the reaction.

The compounds of formula (I-i) wherein Het is an optionally substituted thiazolyl radical, being optionally condensed with a five- or six-membered hetero- or carbocyclic ring, may be prepared by a number of cyclization reactions, yielding, depending upon the case, compounds which may be represented by the formula (I-i-2) or (I-i-3).




R^{24} , R^{25} , R^{26} and R^{27} are each independently optional substituents of the said thiazolyl ring, or, where in the compounds of formula (I-i-2) said thiazolyl ring is condensed with a five- or six-membered hetero- or carbocyclic ring, R^{24} and R^{25} taken together may form a bivalent radical of formula G^3 .


Further, where Het is a radical of formula (e-1), said Het may be formed by condensing an intermediate (XXI) with a $>\text{C}=\text{X}^2$ generating agent, e.g. urea, thiourea, 1,1'-carbonylbis[1H-imidazole], lower alkyl carbonohalidate, phosgene, thiophosgene, trichloromethyl carbonohalidate and the like.



The compounds of formula (I-i-4) wherein R^{11} is hydrogen may additionally be prepared by cyclizing an intermediate of formula




15 reacting a reagent (XXIII) with an amine (XXIV).



The reaction of (XXI) with the >C=X^2 generating agent and the cyclization of (XXII) may conveniently be conducted in a suitable solvent such as, for example, an ether, e.g. 1,1-oxybisethane, tetrahydrofuran, an halogenated hydrocarbon, e.g. dichloromethane, trichloromethane, a hydrocarbon, e.g. benzene, methylbenzene, an alcohol, e.g. methanol, ethanol, a ketone, e.g. 2-propanone, 4-methyl-2-pentanone, N,N-dimethylformamide, N,N-dimethylacetamide, or mixtures of such solvents, optionally in the presence of an appropriate base such as, for example, N,N-diethylethanamine, an alkali or earth alkaline metal carbonate or hydrogen carbonate. In order to enhance the reaction rate, it may be suitable to heat the reaction mixture.

Further, where Het is a radical of formula (e-2), said Het may be generated by cyclizing an intermediate (XXV) with an acid (XXVI) or a suitable functional derivative thereof, thus preparing a compound of


formula (I-i-5). Alternatively an intermediate (XXVII) may be condensed with an aromatic amino acid or -thioacid of formula (XXVIII), preparing also a compound (I-i-5).




15 The reaction of (XXV) with (XXVI) and of (XXVII) with (XXVIII) may be conducted in a suitable reaction-inert solvent, such as, for example, a hydrocarbon, e.g. benzene, methylbenzene, an alcohol, water. In some instances it may be appropriate to use higher temperatures in order to reduce the reaction time.

20 Where Het is a radical of formula (e-3), wherein R^{14} is hydrogen and R^{15} is a radical of formula $R^{15-a}-CH_2-$, said Het may be formed by reacting a compound (XXIX) with an appropriate acetylene derivative (XXX), thus preparing a compound of formula (I-i-6).

25 Additionally, where Het is a radical of formula (e-3), said Het may be formed by reacting (XXIX) with a ketone of formula (XXXI), thus preparing a compound of formula (I-i-7).






5

The reaction of (XXIX) with (XXX) may be conducted in a suitable solvent such as, for example, an alcohol, e.g. methanol, ethanol, while the reaction of (XXIX) with (XXXI) may be conducted in a suitable solvent preferably in the presence of an organic acid such as, for example, ethanedioic acid and the like. Elevated temperatures may also be appropriate to shorten the reaction time.

Additionally, where Het is a radical (e-4), said Het may be created by condensing a reagent (XXXII) with an intermediate (XXXIII), thus giving a compound (I-i-8).


15



20


Where Het is a radical (e-5) being connected to K by the G^3 containing ring and bearing a 2-mercaptosubstituent, said Het may be formed during the cyclization of an intermediate (XXXII) with CS_2 , thus preparing a compound (I-i-9).

25



Where Het is a radical of formula (e-6) being connected to K either by the G^4 containing ring or by the imidazole ring, said Het is formed during the condensation reaction of a reagent (XXXV) with an intermediate (XXXVI) respectively by the cyclodesulfurization reaction of an intermediate (XXXVII), thus preparing a compound (I-i-10) respectively (I-i-11).

35




The reactions of (XXXII) with (XXXIII), of (XXXIV) with CS_2 and (XXXV) with (XXXVI) may conveniently be conducted in a suitable reaction-inert solvent, such as for example one of the solvents given hereinabove for the preparation of (I-i-4) optionally in the presence of an appropriate base, e.g. one of the bases also described for the preparation of (I-i-4); higher temperatures may be used to enhance the reaction rate.

The cyclodesulfurization of (XXXVII) may be carried out by the reaction of (XXXVII) with an appropriate alkyl halide, preferably iodomethane in an appropriate reaction-inert organic solvent, e.g., a lower alkanol such as methanol, ethanol, 2-propanol and the like. Otherwise, the cyclodesulfurization reaction may be carried out by the reaction of (XXXVII) with an appropriate metal oxide or salt in an appropriate solvent according to art-known procedures. For example, the compounds of formula (I) can easily be prepared by the reaction of (XXXVII) with an appropriate Hg(II) or Pb(II) oxide or salt, such as, for example HgO , HgCl_2 , $\text{Hg}(\text{OAc})_2$, PbO or $\text{Pb}(\text{OAc})_2$. In certain instances it may be appropriate to supplement the reaction mixture with a small amount of sulfur. Even so methanediimines, especially N,N'-methanetetraylbis[cyclohexanamine] may be used as cyclodesulfurizing agents.

Where Het is a radical (e-7), said Het may be formed during the condensation of an intermediate (XXXVIII) with a $>C=X^2$ generating agent, following the same procedures as previously described for the preparation of (I-i-4) starting from (XXXIII).

5



10

The compounds of formula (I) can also be converted into each other following art-known procedures of functional group transformation. Some examples will be cited hereinafter.

The compounds of formula (I), wherein -B- is -S- may be converted into the corresponding compounds of formula (I), wherein -B- is -SO- or -SO₂- by an appropriate oxidation reaction, e.g. by reacting the former compounds with a suitable oxidating agent such as, for example, potassium periodate, a peroxide, e.g. 3-chlorobenzenecarboxoperoxoic acid, hydrogen peroxide, and the like, in a suitable solvent such as, for example, an ether, e.g. tetrahydrofuran, 1,1'-oxybisethane, a hydrocarbon, e.g. benzene, a halogenated hydrocarbon, e.g. dichloromethane, trichloromethane and the like. In the instance where a sulfinyl is desired, said oxidation reaction is preferably conducted at lower temperatures with approximately one equivalent of the oxidating agent, while where a sulfonyl is desired, said oxidation reaction may be conducted at room or elevated temperature with an excess of oxidating agent.

The compounds of formula (I) having a nitro substituent can be converted into the corresponding amines by stirring and, if desired, heating the starting nitro-compounds in a hydrogen-containing medium in the presence of a suitable amount of an appropriate catalyst such as, for example, platinum-on-charcoal, palladium-on-charcoal, Raney-nickel and the like catalysts.

Suitable solvents are, for example, alcohols, e.g., methanol, ethanol and the like.

In an analogous procedure, the compounds of formula (I) having a cyano substituent, can be converted into the corresponding aminomethyl containing compounds.

The compounds of formula (I) having an hydroxy substituent may be
5 converted into the corresponding halo compounds following art-known halogenating procedures, e.g., by reacting the former compounds with a suitable halogenating agent, e.g. thionyl chloride, phosphoryl chloride, phosphor trichloride, phosphor pentachloride, thionyl bromide, phosphor tribromide and the like.

10 The compounds of formula (I) containing an ester group may be converted into the corresponding carboxylic acids following art-known saponification procedures, e.g. by treating the said compounds with an aqueous alkaline solution or with an aqueous acidic solution.

The compounds of formula (I) containing a Het substituted with a
15 thioxo group can be converted into the corresponding oxo compounds following art-known procedures, for example, by treating the said thioxo containing compounds with a peroxide, e.g. hydrogen peroxide in a suitable alkaline medium, e.g. an aqueous alkali metal hydroxide solution which may be mixed with an organic solvent, such as, for
20 example, methanol, ethanol and the like.

The compounds of formula (I) containing an unsaturated Het can be converted into the corresponding saturated form following art-known reducing procedures, e.g. by treating the said compounds with hydrogen in the presence of a suitable catalyst such as, for example,
25 platinum-on-charcoal, palladium-on-charcoal and the like catalysts.

Halo atoms substituted on aryl groups may be replaced by hydrogen following art-known hydrogenolysis procedures, i.e. by stirring and, if desired, heating the starting compounds in a suitable solvent under hydrogen atmosphere in the presence of an appropriate catalyst, e.g.,
30 palladium-on-charcoal and the like catalysts. Said halo atoms may also be replaced by a lower alkyloxy or a lower alkylthio substituent by reacting the starting halo-compound with an appropriate alcohol or thioalcohol or, preferably, an alkali- or earth alkaline metal salt or an appropriate alcohol or thioalcohol in a suitable solvent.

35

Lower alkyloxy and lower alkylthio radicals substituted on aryl may be converted into the corresponding hydroxy or thiol radicals by treating them with an aqueous acidic solution, e.g. an aqueous hydrochloric or hydrobromic solution.

- 5 The compounds of formula (I) containing an imino group, e.g. where NR^1 , NR^3 , NR^4 or NR^5 is NH, or an amino group, e.g. where AR^1 , AR^2 or Het is substituted with an amino group, the hydrogen atom in said imino or amino may be replaced by a suitable substituent following art-known procedures such as, for example, N-alkylation, 10 reductive N-alkylation, acylation and the like methods. A number of such procedures will be described hereinafter in more detail. For example, lower alkyl groups or substituted lower alkyl groups may be introduced by reacting the starting compounds with an appropriate N-alkylating agent following the procedures described hereinabove for 15 the N-alkylation reactions of (VII) with (I-c), or by reacting the starting compounds with an appropriate carbonyl-compound following the reductive N-alkylation procedures described hereinabove for the reductive N-alkylations of (I-c-1) with (VIII), (I-d) with (IX) and (I-e) with (X).
- 20 Lower alkylcarbonyl, Ar^2 -carbonyl and the like groups may be introduced by reacting the starting amine with an appropriate carboxylic acid or a derivative thereof such as, for example, an acid halide, acid anhydride and the like.

Lower alkyloxycarbonyl and Ar^2 -oxycarbonyl groups can be 25 introduced by reacting the starting amine compound with an appropriate carbonohalidate, e.g. ethyl carbonohalidate, phenylmethyl carbonohalidate and the like.

Ar^2 -NH-CO, Ar^2 -NH-CS, (lower alkylamino)-CO- (lower alkylamino)-CS-, and the like groups can conveniently be introduced by 30 reacting the starting amine compound with an appropriate isocyanate or isothiocyanate following the procedures described hereinabove for the preparation of (I-b-4), (I-b-5-a), (I-b-5-b) and (I-b-5-c).

The compounds of formula (I) containing a substituted nitrogen atom may be converted into the corresponding compounds of formula (I) 35 wherein said nitrogen bears a hydrogen atom following art-known

methods for preparing N-H groups such as, for example :

1. where said nitrogen is substituted with an $\text{Ar}^2\text{-CH}_2$ group, by treating the starting compounds with hydrogen in the presence of a suitable catalyst, e.g. palladium-on-charcoal, platinum-on-charcoal, in an appropriate solvent;
2. or, where said nitrogen is substituted with a sulfonyl group, e.g. lower alkylsulfonyl and $\text{Ar}^2\text{-sulfonyl}$, by treating the starting compounds with an aqueous acidic solution preferably in the presence of a catalyst such as, for example, phenol, methoxybenzene and the like;
3. or, where said nitrogen atoms are substituted with an $\text{Ar}^2\text{-carbonyl}$ group by treating the starting compounds with an aqueous basic solution, e.g. a alkali metal solution;
4. where said nitrogen is substituted with lower alkyloxy carbonyl or $\text{Ar}^2\text{-oxycarbonyl}$, by treating the starting compounds with an aqueous acidic or aqueous basic solution optionally in admixture with an organic solvent or where said nitrogen atom is substituted with $\text{Ar}^2\text{-oxycarbonyl}$, by catalytically hydrogenating the starting materials in a suitable solvent.
- 20 The compounds of formula (I) containing a nitrogen atom substituted with $\text{Ar}^2\text{-CH}_2\text{-}$ may be converted into the corresponding compounds where said nitrogen is substituted with lower alkyloxycarbonyl, for example by treating the former compounds with a lower alkyl carbonohalidate in the presence of a suitable solvent and, if desired, in the presence of an appropriate base.

The compounds of formula (I) containing a mercapto group may be converted into the corresponding isothiocyanato containing compounds by treating the starting amino compounds with CS_2 in the presence of N,N'-methanetetraylbis[cyclohexanamine].

- 30 The compounds of formula (I) containing a $\text{-CH}_2\text{-C(=O)-}$ fragment can be converted into the corresponding compounds of formula (I) containing a -CH(halo)-C(=O)- fragment following art-known halogenating procedures, e.g. by treating the starting compound with a halogen.

35


In all of the foregoing and in the following preparations, the reaction products may be isolated from the reaction mixture and, if necessary, further purified according to methodologies generally known in the art.

- 5 The compounds of formula (I) have basic properties and, consequently, they may be converted to their therapeutically active non-toxic acid addition salt forms by treatment with appropriate acids, such as, for example, inorganic acids, such as hydrohalic acid, e.g. hydrochloric, hydrobromic and the like, and sulfuric acid, nitric acid, phosphoric acid and the like; or organic acids, such as, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 2-oxopropanoic, ethanedioic, propanedioic, butanedioic, (Z)-2-butenedioic, (E)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxybutanedioic, 2-hydroxy-1,2,3-propanetricarboxylic, 15 methanesulfonic, ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids. Conversely the salt form can be converted by treatment with alkali into the free base form.


- Some intermediates and starting materials in the foregoing 20 preparations are known compounds which may be prepared according to art-known methodologies of preparing said or similar compounds and others are new. A number of such preparation methods will be described hereinafter in more detail.

- The intermediates of formula (II), wherein B is CH_2 , X^1 is NH 25 and W is lower alkyloxy, said intermediates being represented by the formula (II-a), can be prepared by reacting a (cyanomethyl)piperidine of formula (XXXIX) with an alcohol, e.g. methanol, ethanol and the like, in the presence of an acid, e.g. hydrochloric acid.

30




35



5 The intermediates of formula (IV) may be prepared by a reduction reaction of an appropriate 4-piperidinone, and, if desired, followed by an appropriate art-known groupstransformation procedure, e.g., where a compound of formula (V-b) is desired, by reacting the thus obtained alcohol with thionyl chloride, methanesulfonyl chloride and
10 the like in order to obtain an appropriate leaving group.

The intermediates of formula (VI) can be prepared by reacting an appropriate bicyclic condensed imidazole derivative with a pyridinium derivative.



20 The intermediates of formula (VII) can conveniently be prepared following art-known procedures as described in, for example, U.S. Patent Number 4,335,127, U.S. Patent Number 4,342,870 and European Patent Publication Number 0,070,053.

From formula (I) it is evident that the compounds of this
25 invention may have several asymmetric carbon atoms in their structure. Each of these chiral centers may be present in a R- and a S-configuration, this R- and S-notation being in correspondence with the rules described by R.S. Cahn, C. Ingold and V. Prelog in Angew. Chem., Int. Ed. Engl., 5, 385, 511 (1966).

30 Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereoisomers may be separated by physical separation methods such as selective crystallization and chromatographic techniques, e.g., counter current distribution, and enantiomers may be separated from

each other by the selective crystallization of their diastereomeric salts with optically active acids.

Pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically.

It is evident that the cis and trans diastereomeric racemates may be further resolved into their optical isomers, cis(+), cis(-), trans(+) and trans(-) by the application of methodologies known to those skilled in the art.

Stereochemically isomeric forms of the compounds of formula (I) are naturally intended to be embraced within the scope of the invention.

The following examples are intended to illustrate and not to limit the scope of the present invention. Unless otherwise stated all parts therein are by weight.

EXPERIMENTAL PARTA. Preparation of IntermediatesExample 1

5 a) A mixture of 302 parts of ethyl 2-[1-(phenylmethyl)-4-piperi-
dinylidene]acetate hydrochloride and 200 parts of glacial acetic
acid was hydrogenated at normal pressure and at a temperature
between 24-36°C, in the presence of 4 parts of platinum oxide. After
the calculated amount of hydrogen was taken up, the catalyst was
10 filtered off and the filtrate was evaporated. The residue was washed
with 1,1'-oxybisethane, alkalized with sodium hydroxide and
extracted with 1,1'-oxybisethane. The extract was dried over
potassium carbonate and evaporated. The oily residue was distilled in
vacuo, yielding 205 parts of the crude oily free base. From this oil
1 part was taken to prepare the hydrochloride salt. It was dissolved
15 in 1,1'-oxybisethane and gaseous hydrogen chloride was introduced
into the solution: a semi-solid salt was precipitated. The solvent
was decanted and the residue was dissolved again in a mixture of 6
parts of ethanol and 4 parts of 1,1'-oxybisethane. This solution was
concentrated to 5 parts. To the residue were added 12 parts of
20 1,1'-oxybisethane, whereupon a solid was precipitated. It was
filtered off and dried, yielding 0.2 parts of ethyl 1-(phenyl-
methyl)-4-piperidineacetate hydrochloride; mp. 122.5 - 137°C
(intermediate 1)

25 b) A mixture of 8 parts of ethyl 1-(phenylmethyl)-4-piperidine-
acetate hydrochloride and 80 parts of a dilute hydrochloric acid
solution was stirred and refluxed for 4 hours. After cooling, the
reaction mixture was evaporated. The residue was washed with
2-propanone and the solvent was evaporated again. The solid residue
was washed with 2-propanone, filtered off and dried, yielding 6
30 parts of 1-(phenylmethyl)-4-piperidineacetic acid hydrochloride;
mp. 137-145°C (2).

Example 2

To a suspension of 68.5 parts of ethyl 4-oxo-1-piperidine-
carboxylate in 240 parts of methanol were added portionwise 3.8
35 parts of sodium borohydride at a temperature between 20-30°C (the
reaction mixture was cooled if necessary in a water-bath). After the
addition was complete, the whole was stirred for 30 minutes. The

reaction mixture was then poured into a mixture of 53.5 parts of ammonium chloride and 400 parts of water. The methanol was evaporated. The product was extracted with trichloromethane. The extract was dried and evaporated. The oily residue was distilled in vacuo, yielding 60 parts of oily ethyl 4-hydroxy-1-piperidine-carboxylate; bp. 140°C at 1.4 mm. pressure; n_D^{20} : 1.4796; d_4^{20} : 1.1166 (3).

In a similar manner there was also prepared:

methyl (cis+trans)-4-hydroxy-3-methyl-1-piperidinecarboxylate (4).

Example 3

To a stirred solution of 90 parts of 1-[(4-methylphenyl)sulfonyl]-4-piperidinol, 37.5 parts of N,N-diethylethanamine and 1300 parts of dichloromethane was added dropwise a solution of 42.3 parts of methanesulfonyl chloride in 130 parts of dichloromethane (exothermic reaction: temperature rose to 35°C). Upon completion, stirring was continued for 2 hours at room temperature. Water was added and the layers were separated. The organic phase was washed with water, dried, filtered and evaporated. The residue was suspended in 2,2'-oxybispropane. The product was filtered off and dried, yielding 116 parts (100%) of 1-[(4-methylphenyl)sulfonyl]-4-piperidinol methanesulfonate (ester); mp. 168.5 - 175.3°C (5).

Example 4

2350 Parts of hydrogen chloride were bubbled through 5600 parts of cooled ethanol (ice bath) at 10°C. Then there were added dropwise, during a 45 minutes period, 1500 parts of 1-(phenylmethyl)-4-piperidineacetonitrile. Upon completion, the whole was stirred for 20 hours at room temperature. The reaction mixture was evaporated and the residue was stirred in 2400 parts of acetonitrile. The product was filtered off, washed with 560 parts of acetonitrile and dried, yielding 2000 parts (85.7%) of O-ethyl 1-(phenylmethyl)-4-piperidineethanimidate hydrochloride (6)

In a similar manner there was also prepared:

O-methyl 1-(phenylmethyl)-4-piperidineethanimidate dihydrochloride (7).

Example 5

A mixture of 180.0 parts of 2-chloro-3-nitropyridine, 122.0 parts of 2-thiophenemethanamine, 191.0 parts of sodium carbonate, 1 part of potassium iodide and 810 parts of N,N-dimethylacetamide was

stirred for 1.50 hours at 100°C. The reaction mixture was poured into about 4000 parts of water. The whole was stirred overnight at room temperature. The precipitated product was filtered off and dried in vacuo at 40°C, yielding 251.5 parts of 3-nitro-N-(2-thienylmethyl)-2-pyridinamine; mp. 100°C (8).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:

N-(2-nitrophenyl)-3-pyridinemethanamine (9);

N-(4-fluorophenylmethyl)-3-nitro-2-pyridinamine; mp. 76°C (10)

10 N-(3-nitro-2-pyridinyl)-2-pyridinemethanamine; mp. 113.6°C (11)

2-nitro-N-(2-thienylmethyl)benzenamine (12)

4-methyl-N-(2-nitrophenyl)benzenemethanamine; mp. 65°C (13)

N-[(4-methylphenyl)methyl]-3-nitro-2-pyridinamine; mp. 80.0-87.3°C (14).

15 N³-[(4-fluorophenyl)methyl]-2,3-pyridinediamine (15);

N-[(4-fluorophenyl)methyl]-3-nitro-4-pyridinamine; mp. 136.8°C (16);

N-[(4-fluorophenyl)methyl]-4-nitro-3-pyridinamine, 1-oxide (17);

4-fluoro-N-(4-methoxy-2-nitrophenyl)benzenemethanamine (18);

4-fluoro-N-(5-methoxy-2-nitrophenyl)benzenemethanamine (19);

20 4-fluoro-N-(4-methyl-2-nitrophenyl)benzenemethanamine; mp. 99.9°C (20);

4-fluoro-N-(3-methoxy-2-nitrophenyl)benzenemethanamine; (21);

4-fluoro-N-(2-methoxy-6-nitrophenyl)benzenemethanamine; (22);

4-fluoro-N-(4,5-dimethoxy-2-nitrophenyl)benzenemethanamine; (23);

4-fluoro-N-(4-chloro-5-methoxy-2-nitrophenyl)benzenemethanamine; (24);

25 4-fluoro-N-(5-chloro-4-methoxy-2-nitrophenyl)benzenemethanamine; (25);

N-(4-methoxy-2-nitrophenyl)-2-furanmethanamine; (26);

N-(5-methoxy-2-nitrophenyl)-2-furanmethanamine; (27);

N-(4-methoxy-2-nitrophenyl)-2-pyridinemethanamine; (28);

N-(5-methoxy-2-nitrophenyl)-2-pyridinemethanamine; (29);

30 N-[(4-fluorophenyl)methyl]-6-methoxy-3-nitro-2-pyridinamine; (30);

N-[(2-furanyl)methyl]-6-methoxy-3-nitro-2-pyridinamine; (31); and

N-(3-nitro-6-methoxy-2-pyridinyl)-2-pyridinemethanamine; (32).

Example 6

To a stirred and cooled mixture of 40 parts of N-[(4-fluorophenyl)-methyl]-4-nitro-3-pyridinamine, 1-oxide and 1050 parts of trichloromethane were added dropwise 47 parts of phosphor penta-chloride at a

temperature between 0 and -10°C. Upon completion, the whole was stirred and refluxed for 1 hour. While stirring, the reaction mixture was cooled. The precipitated product was filtered off, stirred in water and alkalized with ammonium hydroxide. The product was extracted with trichloromethane. The extract was dried, filtered and vaporated. The residue was stirred in 2,2'-oxybispropane. The product was filtered off and dried, yielding 22.2 parts of N-[(4-fluorophenyl)methyl]-4-nitro-3-pyridinamine; mp. 91.9°C (33).

Example 7

A mixture of 100 parts of N-[(4-methoxyphenyl)methyl]-3-nitro-2-pyridinamine, 3 parts of a solution of thiophene in methanol 4% and 480 parts of methanol saturated with ammonia was hydrogenated at normal pressure and at 50°C with 5 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated, yielding 88.4 parts of N²-[(4-methoxyphenyl)methyl]-2,3-pyridinediamine; mp. 118.1°C (34).

In a similar manner there were also prepared:

N-(3-pyridinylmethyl)-1,2-benzenediamine (35);
N²-[(4-fluorophenyl)methyl]-2,3-pyridinediamine (36);
N²-(2-pyridinylmethyl)-2,3-pyridinediamine; mp. 134.9 (37);
N²-(2-furanylmethyl)-2,3-pyridinediamine (38);
N¹-(2-thienylmethyl)-1,2-benzenediamine (39);
N²-(2-thienylmethyl)-2,3-pyridinediamine; mp. 92.1°C (40);
N¹-[(4-methylphenyl)methyl]-1,2-benzenediamine (41);
N²-[(4-methylphenyl)methyl]-2,3-pyridinediamine; mp. 125.1°C (42);
N⁴-[(4-fluorophenyl)methyl]-3,4-pyridinediamine; mp. 163.7°C (43);
N³-[(4-fluorophenyl)methyl]-3,4-pyridinediamine; mp. 159.6°C (44);
N¹-[(4-fluorophenyl)methyl]-4-methoxy-1,2-benzenediamine (45);
N²-[(4-fluorophenyl)methyl]-4-methoxy-1,2-benzenediamine (46);
N¹-[(4-fluorophenyl)methyl]-4-methyl-1,2-benzenediamine (47);
N-[(5-methyl-2-furanyl)methyl]-1,2-benzenediamine (48);
N¹-[(4-fluorophenyl)methyl]-3-methoxy-1,2-benzenediamine (49);
N¹-[(4-fluorophenyl)methyl]-6-methoxy-1,2-benzenediamine (50);
N¹-[(4-fluorophenyl)methyl]-4,5-dimethoxy-1,2-benzenediamine (51);
N¹-[(4-fluorophenyl)methyl]-5-chloro-4-methoxy-1,2-benzenediamine (52);
N¹-[(4-fluorophenyl)methyl]-4-chloro-5-methoxy-1,2-benzenediamine (53);
N¹-(2-furanylmethyl)-4-methoxy-1,2-benzenediamine (54);

N^1 -(2-furanylmethyl)-5-methoxy-1,2-benzenediamine (55);
 N^1 -(2-pyridinylmethyl)-4-methoxy-1,2-benzenediamine (56);
 N^1 -(2-pyridinylmethyl)-5-methoxy-1,2-benzenediamine (57);
 N^2 -[(4-fluorophenyl)methyl]-6-methoxy-2,3-pyridinediamine (58);
5 N^2 -(2-furanylmethyl)-6-methoxy-2,3-pyridinediamine (59); and
 N^2 -(2-pyridinylmethyl)-6-methoxy-2,3-pyridinediamine (60);

Example 8

A mixture of 60 parts of 2-chloro-1H-benzimidazole, 58 parts of
1-(chloromethyl)-4-fluorobenzene, 42.5 parts of sodium carbonate, 0.1
10 parts of potassium iodide and 135 parts of N,N-dimethyl-formamide was
stirred and heated overnight at 70°C. The reaction mixture was poured
into water. The precipitated product was filtered off and dissolved in
trichloromethane. The solution was dried, filtered and evaporated. The
residue was crystallized from 2,2'-oxybispropane, yielding 62.5 parts of
15 2-chloro-1-(4-fluorophenylmethyl)-1H-benzimidazole (61).

Example 9

A mixture of 8.35 parts of thiourea, 26 parts of 2-chloro-1-
[(4-fluorophenyl)methyl]-1H-benzimidazole and 400 parts of ethanol was
stirred and refluxed for 5 hours. The reaction mixture was evaporated.
20 The residue was suspended in 2,2'-oxybispropane. The precipitated
product was filtered off and crystallized from ethanol, yielding 6.1
parts of 1-[(4-fluorophenyl)methyl]-1H-benzimidazole-2-thiol; mp.
194.7°C (62).

Example 10

25 To a stirred solution of 6 parts of 1,2-dimethyl-1H-benzimidazole in
50 parts of dry pyridine were added dropwise 6.2 parts of benzoyl
chloride at room temperature. Upon completion, stirring was continued
for 2 hours at room temperature. The whole was evaporated. The residue
was dissolved in 260 parts of dichloromethane. Water was added and the
30 solution was treated with concentrate ammonium hydroxide. The dichloro-
methane layer was decanted, dried, filtered and evaporated. The residue
was taken up twice in 45 parts of methylbenzene and the latter was
evaporated each time, yielding 1-benzoyl-1,4-dihydro-4-[(1-methyl-
1H-benzimidazol-2-yl)methyl]pyridine as an oily residue (63).

35 In a similar manner there was also prepared:
ethyl 4-[(1-methyl-1H-benzimidazol-2-yl)methyl]-1(4H)-pyridine-
carboxylate as an oily residue (64).

Example 11

A mixture of 9.7 parts of 4-fluoro- δ -(4-fluorophenyl)benzen - butanoyl chloride, 4.1 parts of 2,6-dimethylpyridine and 68 parts of tetrahydrofuran was hydrogenated at normal pressure and at room temperature with 2 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated, yielding 8.5 parts of 4-fluoro- δ -(4-fluorophenyl)benzenebutanal (65).

Example 12

To a stirred mixture of 26 parts of 1-ethyl-1,4-dihydro-5H-tetrazole-5-thione, 13.8 parts of potassium carbonate and 240 parts of 2-propanone were added dropwise 376 parts of 1,2-dibromoethane. Upon completion, stirring was continued overnight. The precipitate was filtered off and the filtrate was evaporated, yielding 45 parts (95%) of 5-[(2-bromoethyl)thio]-1-ethyl-1H-tetrazole as a residue (66).

Example 13

To a stirred and cooled (0-10°C) mixture of 59 parts of 2-propanamine and 650 parts of dichloromethane were added dropwise 94.2 parts of 3-bromopropanoyl chloride. Upon completion, stirring was continued for 5 minutes. The whole was washed with water. The organic layer was separated, dried, filtered and evaporated. The residue was crystallized from a mixture of 2,2'-oxybispropane and hexane. The product was filtered off and dried, yielding 70 parts (73%) of 3-bromo-N-(1-methylethyl)propanamide; mp. 60°C (67).

Example 14

A mixture of 4.76 parts of 6-chloro-N⁴-methyl-4,5-pyridinediamine, 26.6 parts of 1,1,1-triethoxyethane and 30 parts of acetic acid anhydride was stirred and refluxed for 3 hours. The reaction mixture was evaporated. The residue was crystallized from a mixture of hexane and methylbenzene. The product was filtered off and dried, yielding 5.3 parts (96.3%) of 6-chloro-8,9-dimethyl-9H-purine (68).

Example 15

A mixture of 4.76 parts of 6-chloro-N⁴-methyl-4,5-pyrimidinediamine and 7.2 parts of urea was stirred and heated for 1 hour at 180°C. After cooling, the residue was suspended in water. The product was filtered off and dried, yielding 3.3 parts (60%) of 6-chloro-9-methyl-9H-purin-8-ol (69).

B. Preparation of Final CompoundsExample 16

To 73 parts of hot (70°C) polyphosphoric acid were added 27 parts of 1-(phenylmethyl)-4-piperidineacetic acid hydrochloride: temperature rose to 100°C. When the addition was complete, there were added portionwise 14 parts of 1,2-benzenediamine and stirring and heating was continued for 50 minutes at 170°C. The hot reaction mixture was poured into 300 parts of warm water. The acid solution was alkalinized with a potassium hydroxide solution. The precipitated free base was filtered off, washed with water and extracted with trichloromethane. The extract was dried and evaporated. The solid residue was recrystallized from a mixture of 2-propanone and methanol, yielding 17 parts of 2-[[1-(phenylmethyl)-4-piperidinyl]-methyl]-1H-benzimidazole; mp. 221.5-222°C (compound 1).

In a similar manner there was also prepared:
2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-imidazo[4,5-c]pyridine; mp. 172.9°C (2).

Example 17

A mixture of 27.3 parts of O-methyl 1-(phenylmethyl)-4-piperidineethanimidate dihydrochloride, 14 parts of N-(2-furanylmethyl)-1,2-benzenediamine and 250 parts of acetic acid was stirred overnight at room temperature. The reaction mixture was evaporated and water was added to the residue. The whole was alkalinized with sodium carbonate and extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was crystallized from 1,1'-oxybisethane. The product was filtered off and dried, yielding 15.5 parts (57%) of 1-(2-furanylmethyl)-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole; mp. 124.8°C (3).

In a similar manner there were also prepared:
1-phenyl-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole; mp. 141.6°C (4); and
2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1-(3-pyridinylmethyl)-1H-benzimidazole; mp. 125.4°C (5).

Example 18

A mixture of 116.5 parts of O-ethyl 1-(phenylmethyl)-4-piperidineethanimidate hydrochloride, 61.5 parts of N¹-[(4-methylphenyl)methyl]-1,2-benzenediamine and 400 parts of methanol was stirred and refluxed overnight. An ther portion of 40 parts of O-ethyl 1-(phenylmethyl)-4-piperidineethanimidate hydrochloride was add d and stirring was continued for 4 hours at reflux. The reaction mixture was evaporated. Water was added to the residue. The solution was treated with ammonium hydroxide. The product was extracted with
 0 trichloromethane. The extract was dried, filtered and evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 74.5 parts (63%) of 1-[(4-methylphenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole; mp. 124.2°C (6).

.5 Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:

- 2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1-(2-thienylmethyl)-1H-benzimidazole; mp. 156.3°C (7);
- 3-[(4-fluorophenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]-
 20 methyl]-3H-imidazo[4,5-b]pyridine; mp. 103.2-105.8°C (8);
- 2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-3-(2-pyridinylmethyl)-3H-imidazo[4,5-b]pyridine; mp. 118.5-120.9°C (9);
- 3-(2-furanylmethyl)-2-[[1-(phenylmethyl)-4-piperidinyl]-methyl]-3H-imidazo[4,5-b]pyridine; mp. 118.5-119.8°C (10);
- 25 1-[(4-methoxyphenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]-methyl]-1H-benzimidazole; mp. 95.2°C (11);
- 2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-3-(2-thienylmethyl)-3H-imidazo[4,5-b]pyridine; mp. 115.2°C (12);
- 3-(phenylmethyl)-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-3H-
 30 imidazo[4,5-b]pyridine (13);
- 1-(phenylmethyl)-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole; mp. 130°C (14);
- 3-[(4-methylphenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]-methyl]-3H-imidazo[4,5-b]pyridine (15);
- 35 3-[(4-methoxyphenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]-

- methyl]-3H-imidazo[4,5-b]pyridine; mp. 83.4°C (16);
1-[(4-fluorophenyl)methyl]-5-methoxy-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole; mp. 112.6°C (17);
1-(3-furanylmethyl)-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole; mp. 102.0°C (18);
1-[(4-fluorophenyl)methyl]-5-methyl-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole (19);
1-[(4-fluorophenyl)methyl]-6-methoxy-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole; mp. 110°C (20);
5-fluoro-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole; mp. 206.2°C (21); and
1-[(5-methyl-2-furanyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole; mp. 96°C (22).

Example 19

- 15 A mixture of 43 parts of 1-(phenylmethyl)-4-piperidineacetic acid hydrochloride, 31.5 parts of N^3 -[(4-fluorophenyl)methyl]-2,3-pyridinediamine, 850 parts of phosphoryl chloride and 20 parts of N,N-diethylbenzenamine was stirred for 6 hours at reflux temperature. The reaction mixture was evaporated. Methylbenzene was
20 added twice to the residue and the whole was each time evaporated. The final residue was poured into ice water and the whole was made alkaline with a dilute sodium hydroxide solution. The product was extracted twice with dichloromethane. The combined extracts were washed twice with water, dried, filtered and evaporated. The residue
25 was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 30
30 parts (50.2%) of 1-[(4-fluorophenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-imidazo[4,5-b]pyridine; mp. 140.7°C (23).

In a similar manner there were also prepared:

- 1-[(4-fluorophenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-imidazo[4,5-c]pyridine; mp. 139.1°C (24); and
35 3-[(4-fluorophenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]-

methyl]-3H-imidazo[4,5-c]pyridine; mp. 116.9°C (25).

Example 20

To a stirred mixture of 3.5 parts of ethyl 4-hydroxy-1-piperidinecarboxylate and 135 parts of N,N-dimethylformamide was added 1 part of a sodium hydride dispersion 50% and stirring was continued for 2 hours at room temperature. After the addition of 5.2 parts of 2-chloro-1-[(4-fluorophenyl)methyl]-1H-benzimidazole, the whole was further stirred overnight at room temperature. The reaction mixture was poured into ice water and the product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was crystallized from 2,2'-oxybispropane, yielding 2.5 parts (31.5%) of ethyl 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]oxy]-1-piperidinecarboxylate; mp. 94.0°C (26).

In a similar manner there was also prepared:

methyl (cis+trans)-4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]oxy]-3-methyl-1-piperidinecarboxylate (27).

Example 21

To a stirred mixture of 1.5 parts of 1H-benzimidazole-2-thiol and 90 parts of N,N-dimethylformamide were added 0.8 parts of a sodium hydride dispersion 50%. Stirring was continued for 1 hour. Then there were added 3.3 parts of 1-[(4-methylphenyl)sulfonyl]-4-piperidinol methanesulfonate(ester) and the whole was stirred overnight at room temperature. Stirring was continued overnight at 80°C. The reaction mixture was poured into water. The product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was crystallized from acetonitrile, yielding 0.7 parts (18%) of 4-[[1H-benzimidazol-2-yl]thio]-1-[(4-methylphenyl)sulfonyl]piperidine; mp. 194.8°C (28).

In a similar manner there was also prepared:

4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]thio]-1-[(4-methylphenyl)sulfonyl]piperidine; mp. 167.2°C (29).

Example 22

To a stirred and cooled (0°C) mixture of 7.2 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]thio]-1-[(4-

methylphenyl)sulfonyl]piperidine and 95 parts of dichloromethane was added dropwise a solution of 2.6 parts of 3-chlorobenzenecarboxylic acid in dichloromethane. Upon completion, stirring was continued for 2 hours at room temperature. The reaction mixture was washed with a sodium carbonate solution and with water. The organic layer was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 0.7 parts (9%) of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]sulfonyl]-1-[(4-methylphenyl)sulfonyl]piperidine; mp. 166.2°C (30).

Example 23

To a stirred solution of 7.2 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]thio]-1-[(4-methylphenyl)sulfonyl]piperidine in 195 parts of dichloromethane was added dropwise a solution of 7 parts of 3-chlorobenzenecarboxylic acid in 65 parts of dichloromethane. Upon completion, stirring was continued for 2 hours at room temperature. The whole was washed with a sodium carbonate solution and twice with water, dried, filtered and evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 3 parts (40%) of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]sulfonyl]-1-[(4-methylphenyl)sulfonyl]piperidine; mp. 214.7°C (31).

Example 24

A mixture of 16 parts of 1-benzoyl-1,4-dihydro-4-[(1-methyl-1H-benzimidazol-2-yl)methyl]pyridine and 160 parts of methanol was hydrogenated at normal pressure and at 50°C with 5 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (97.5:2.5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The oily residue was

crystallized from 14 parts of 1,1'-oxybis thane. The product was filter d off and dried, yielding 7.8 parts (58.5%) of 1-benzoyl-4-[(1-methyl-1H-benzimidazol-2-yl)methyl]pyridine; mp. 159.3°C (32).

In a similar manner there was also prepared:

- 5 ethyl 4-[(1-methyl-1H-benzimidazol-2-yl)methyl]-1-piperidine-carboxylate; mp. 98.2°C (33).

Example 25

- To a stirred mixture of 55 parts of 2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole and 450 parts of 10 N,N-dimethylformamide were added 10.6 parts of a sodium hydride dispersion 50% and stirring was continued for 1 hour. Then there were added dropwise 26 parts of 1-(chloromethyl)-4-fluorobenzene (slightly exothermic reaction). Upon completion, stirring was continued overnight at room temperature. Water was added and the 15 product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The solid residue was stirred in 1,1'-oxybisethane. The product was filtered off and dried, yielding 67.6 parts (90%) of 1-[(4-fluorophenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole; mp. 127.5°C (34).

- 20 Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:

2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1-(4-thiazolylmethyl)-1H-benzimidazole; mp. 98.7-105.8°C (35); and

- 5(or 6)-fluoro-1-[(4-fluorophenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole (36)

Example 26

- A mixture of 41 parts of 3-[(4-methoxyphenyl)methyl]-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-3H-imidazo[4,5-b]-pyridine and 480 parts of methanol was hydrogenated at normal 30 pressure and at 50°C with 5 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated, yielding 30 parts (89%) of 3-[(4-methoxyphenyl)methyl]-2-(4-piperidinylmethyl)-3H-imidazo[4,5-b]pyridine (37).

- 35 In a similar manner there were also prepared:

- 2-(4-piperidinylmethyl)-1H-benzimidazole; mp. 195-196.5°C (38);
3-[(4-methylphenyl)methyl]-2-(4-piperidinylmethyl)-3H-imidazo-
[4,5-b]pyridine (39);
1-(phenylmethyl)-2-(4-piperidinylmethyl)-1H-benzimidazole.
5 monohydrate; mp. 72.5°C (40);
3-(phenylmethyl)-2-(4-piperidinylmethyl)-3H-imidazo[4,5-b]pyridine
(41);
1-[(4-methylphenyl)methyl]-2-(4-piperidinylmethyl)-1H-benzimidazole
ethanedioate (1:2). monohydrate; mp. 195.1°C (42);
10 1-[(4-methoxyphenyl)methyl]-2-(4-piperidinylmethyl)-1H-benzimidazole
ethanedioate (1:2). monohydrate; mp. 172.1°C (43);
2-(4-piperidinylmethyl)-3-(2-pyridinylmethyl)-3H-imidazo[4,5-b]-
pyridine (E)-2-butenedioate (2:3); mp. 191.1-194.0°C (44);
3-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-3H-imidazo[4,5-b]-
15 pyridine (E)-2-butenedioate (1:2); mp. 200.0-201.2°C (45);
1-phenyl-2-(4-piperidinylmethyl)-1H-benzimidazole; mp. 142.6°C (46);
1-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-1H-benzimidazole
(E)-2-butenedioate (2:3); mp. 204.7°C (47);
1-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-1H-imidazo[4,5-b]-
20 pyridine (E)-2-butenedioate(2:5); mp. 214.4°C (48);
1-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-1H-imidazo[4,5-c]-
pyridineethanedioate(2:3).monohydrate; mp. 173.5°C (49);
3-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-3H-imidazo[4,5-c]-
pyridine (E)-2-butenedioate(2:5); mp. 168.8°C (50);
25 1-[(4-fluorophenyl)methyl]-5-methoxy-2-(4-piperidinylmethyl)-1H-benz-
imidazole dihydrochloride.monohydrate; mp. 214.1°C (51);
1-[(4-fluorophenyl)methyl]-5-methyl-2-(4-piperidinylmethyl)-1H-benz-
imidazole (52); and
1-[(4-fluorophenyl)methyl]-6-methoxy-2-(4-piperidinylmethyl)-1H-benzim-
30 idazole (53).
5(or 6)-fluoro-1-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-1H-
benzimidazole (54).

Example 27

- A mixture of 4.95 parts of 4-[[1-[(4-fluorophenyl)methyl]-
35 1H-benzimidazol-2-yl]thio]-1-[(4-methylphenyl)sulfonyl]piperidine,

225 parts of a hydrobromic acid solution 48% in water and 5 parts of phenol was stirred and refluxed for 2 hours. The reaction mixture was evaporated and the residue was taken up in water and treated with a sodium hydroxide solution. The product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was purified by filtration over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated, yielding 3.4 parts (99%) of 1-[(4-fluorophenyl)-methyl]-2-[(4-piperidinyl)thio]-1H-benzimidazole (55).

Example 28

A mixture of 3.3 parts of 1-benzoyl-4-[(1-methyl-1H-benzimidazol-2-yl)methyl]piperidine, 1.7 parts of water and 40 parts of 2-propanol was stirred and refluxed for 30 hours. The reaction mixture was concentrated and the residue was shaken with 260 parts of dichloromethane. The formed precipitate was filtered off and the filtrate was washed thoroughly with 20 parts of water. The organic phase was dried, filtered and evaporated. The residue was converted into the ethanedioate salt in ethanol. The salt was filtered off and dried, yielding 3.4 parts (83%) of 1-methyl-2-(4-piperidinylmethyl)-1H-benzimidazole ethanedioate (1:2); mp. 219.7°C (56).

Example 29

To a stirred mixture of 76 parts of 2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1-(3-pyridinylmethyl)-1H-benzimidazole and 360 parts of methylbenzene were added dropwise 41 parts of ethyl carbonochloridate. Upon completion, stirring was continued for 2 hours at reflux. Another portion of 5 parts of ethyl carbonochloridate was added dropwise. Upon completion, stirring was continued for 2 hours at reflux. After cooling, the organic layer was washed with a sodium carbonate solution, dried, filtered and evaporated. The residue was purified by filtration over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated, yielding 35.6 parts (50%) of ethyl 4-[[1-(3-pyridinyl)-methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinecarboxylate (57).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:

- methyl 4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinecarboxylate (58);
- 5 ethyl 4-[[1-(2-thienylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinecarboxylate monohydrochloride; mp. 178.7°C (59);
- ethyl 4-[[1-(4-thiazolylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinecarboxylate monohydrochloride; mp. 197.4-199.2°C (60);
- ethyl 4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-
- 10 piperidinecarboxylate as a residue (61);
- ethyl 4-[[3-(2-thienylmethyl)-3H-imidazo[4,5-b]-2-yl]methyl]-1-piperidinecarboxylate as a residue (62); and
- ethyl 4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinecarboxylate (63).

15 Example 30

- A mixture of 68 parts of ethyl 4-[[1-(2-thienylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinecarboxylate monohydrochloride, 95 parts of potassium hydroxide, 800 parts of 2-propanol and 10 parts of water was stirred and refluxed for 6 hours. The
- 20 reaction mixture was evaporated and water was added to the residue. The product was extracted with methylbenzene. The extract was dried, filtered and evaporated. The residue was crystallized from 1,1'-oxybisethane. The product was filtered off and dried, yielding 27 parts (49%) of 2-(4-piperidinylmethyl)-1-(2-thienylmethyl)-1H-benz-
- 25 imidazole; mp. 117.4°C (64).

In a similar manner there were also prepared:

- 1-(2-furanylmethyl)-2-(4-piperidinylmethyl)-1H-benzimidazole
- (E)-2-butenedioate (2:3); mp. 219.6°C (65);
- 1-[(4-fluorophenyl)methyl]-2-(4-piperidinyl)-1H-benzimidazole
- 30 dihydrochloride; mp. 145.2°C (66)
- 2-(4-piperidinylmethyl)-1-(3-pyridinylmethyl)-1H-benzimidazole as a residue (67);
- 3-(2-furanylmethyl)-2-(4-piperidinylmethyl)-3H-imidazo[4,5-b]pyridine
- ethanedioate (2:3). monohydrate; mp. 136.7°C (68);
- 35 2-(4-piperidinylmethyl)-3-(2-thienylmethyl)-3H-imidazo[4,5-b]-

pyridine (E)-2-butenedioate (2:3); mp. 209.6°C (69);

cis-1-[(4-fluorophenyl)methyl]-2-[(3-methyl-4-piperidinyl)oxy]-1H-benzimidazole monohydrochloride monohydrate mp. 143.7°C (70); and

trans-1-[(4-fluorophenyl)methyl]-2-[(3-methyl-4-piperidinyl)oxy]-1H-benzimidazole dihydrochloride; mp. 111.6°C (71).

Example 31

A mixture of 2 parts of ethyl 4-[[1-(4-thiazolylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinecarboxylate and 30 parts of a hydrobromic acid solution 48% was stirred and refluxed for 15 minutes. The reaction mixture was evaporated. The oily residue was crystallized from a mixture of ethanol and 2-propanone, yielding 2 parts of 2-(4-piperidinylmethyl)-1-(4-thiazolylmethyl)-1H-benzimidazole trihydrobromide; mp. 208.3-226.3°C (72).

Example 32

To a stirred mixture of 72 parts of 1-(3-furanylmethyl)-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1H-benzimidazole and 324 parts of methylbenzene were added dropwise 25.5 parts of ethyl carbonochloridate at reflux. Upon completion, stirring was continued for 2 hours at reflux temperature. After cooling, the mixture was washed twice with a sodium hydroxide solution 5%, once with water, dried, filtered and evaporated. This residue, together with 560 parts of 2-propanol, 69.9 parts of potassium hydroxide and 6 parts of water, was stirred and refluxed for 22 hours. The whole was cooled and evaporated. The residue was taken up in water. The product was extracted three times with dichloromethane. The combined extracts were washed twice with water, dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane, methanol and ammonium hydroxide (90:10:1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was stirred in 2-propanone. The product was filtered off and dried, yielding 28.0 parts (49.8%) of 1-(3-furanylmethyl)-2-(4-piperidinylmethyl)-1H-benzimidazole; mp. 180°C (73).

In a similar manner there was also prepared:

1-[(5-methyl-2-furanyl)methyl]-2-(4-piperidinylmethyl)-1H-benzimidazole; mp. 90°C (74).

Example 33

5 T a stirred mixture of 10.2 parts of 1-[(4-fluorophenyl)methyl]-2-[(4-piperidinylthio)-1H-benzimidazole, 3.1 parts of N,N-diethylethanamine and 130 parts of dichloromethane was added dropwise a solution of 5.12 parts of (phenylmethyl) carbonochloridate in 65 parts of dichloromethane. Upon completion, stirring was continued for 1 hour at room temperature. The reaction mixture was washed with
10 water. The organic layer was dried, filtered and evaporated, yielding 14.3 parts of (phenylmethyl) 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]thio]-1-piperidinecarboxylate as a residue (75).

In a similar manner there were also prepared:

15 (phenylmethyl) 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]sulfonyl]-1-piperidinecarboxylate.; mp. 147.3°C (76); and 1-[(4-fluorophenyl)methyl]-2-(4-piperidinylsulfonyl)-1H-benzimidazole.; mp. 133.5°C (77).

Example 34

20 A solution of 22.3 parts of 1-[(4-fluorophenyl)methyl]-5-methoxy-2-(4-piperidinylmethyl)-1H-benzimidazole dihydrochloride monohydrate in 75 parts of a hydrobromic acid solution 48% in water was stirred and refluxed for 18 hours. The whole was slightly cooled and evaporated. The residue was dissolved in water. The solution was
25 treated with an ammonium hydroxide solution. The product was extracted three times with trichloromethane. The combined organic layers were washed with water, dried, filtered and evaporated, yielding 15.7 parts (92%) of 1-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-1H-benzimidazol-5-ol; mp. 210°C (78).

30 In a similar manner there was also prepared: 1-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-1H-benzimidazol-6-ol (79).

Example 35

35 To a stirred mixture of 19.9 parts of 1-[(4-fluorophenyl)methyl]-2-(4-piperidinylloxy)-1H-benzimidazole dihydrochloride, 12.2

parts of N,N-diethylethanamine and 65 parts of dichloromethane was added a solution of 6.5 parts of 2-furanacetic acid and 20.6 parts of N,N-methanetetraylbis[cyclohexanamine] in 130 parts of dichloromethane. The whole was stirred over weekend at room temperature. The reaction mixture was filtered and the filtrate was poured into water. The organic phase was separated, dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from 1,1'-oxybisethane. The product was filtered off and dried, yielding 5.3 parts (25%) of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]oxy]-1-[2-(2-fuanyl)acetyl]piperidine; mp. 128.7°C (80).


Example 36

A mixture of 2 parts of poly(oxymethylene), 3.5 parts of 1-[(4-methylphenyl)methyl]-2-(4-piperidinylmethyl)-1H-benzimidazole, 1 part of a solution of thiophene in ethanol 4% and 120 parts of methanol was hydrogenated at normal pressure and at 50°C with 2 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated. The residue was taken up in water. The product was extracted with dichloromethane. The organic layer was dried, filtered and evaporated. The residue was converted into the (E)-2-butenedioate salt in ethanol. The salt was filtered off and dried, yielding 3 parts (536%) of 1-[(4-methylphenyl)methyl]-2-[[1-methyl-4-piperidinyl)methyl]-1H-benzimidazole (E)-2-butenedioate (2:3); mp. 188.9°C (81).

30

35

In a similar manner there were also prepared:



5

No.	L	R ¹	A ¹	B	salt/ base	mp. °C
10	82 CH ₃	4-F-C ₆ H ₄ CH ₂	CH	CH ₂	*	193.1
	83 (CH ₃) ₂ CH	4-F-C ₆ H ₄ CH ₂	CH	CH ₂	*	165.7
	84 (CH ₃) ₂ CH	4-F-C ₆ H ₄ CH ₂	CH	O	*	210.5
	85 CH ₃	4-F-C ₆ H ₄ CH ₂	CH	O	*	161.0
	86 CH ₃	2-furanylmethyl	CH	CH ₂	*	178.4
15	87 (CH ₃) ₂ CH	2-furanylmethyl	CH	CH ₂	HCl	206.0
	88 CH ₃	4-thiazolylmethyl	CH	CH ₂	**	146.9
	89 (CH ₃) ₂ CH	4-F-C ₆ H ₄ CH ₂	N	CH ₂	***	163.0
	90 CH ₃	4-F-C ₆ H ₄ CH ₂	N	CH ₂	***	158.2
	91 CH ₃	H	CH	CH ₂	base oil	
20	92 (4-F-C ₆ H ₄) ₂ CH(CH ₂) ₃	H	CH	CH ₂	***	212.5
	93 CH ₃	C ₆ H ₅ CH ₂	CH	CH ₂	base	96.1
	94 (CH ₃) ₂ CH	C ₆ H ₅ CH ₂	CH	CH ₂	***	194.2
	95 (CH ₃) ₂ CH	CH ₃	CH	CH ₂	***	116.3
	96 CH ₃	4-F-C ₆ H ₄ CH ₂	CH	S	***	139.0
25	97 C ₆ H ₅ CH ₂ -N	4-F-C ₆ H ₄ CH ₂	CH	CH ₂	2HCl	>300
				H ₂ O		(dec.)
	98 cyclohexyl	4-F-C ₆ H ₄ CH ₂	N	CH ₂	***	163.9
	99 cyclohexyl	4-F-C ₆ H ₄ CH ₂	CH	CH ₂	***	172.1

30 * : (E)-2-butenedioate salt (2:3)

** : ethanedioate (1:2) salt.monohydrate

***: ethanedioate (1:2)

In a similar manner there was also prepared:

1-[(4-fluorophenyl)methyl]-2-[[1-(1-methylethyl)-4-piperidinyl]-
 35 methyl]-1H-imidazo[4,5-c]pyridine; mp. 115.9°C (100).

Example 37

A mixture of 7.9 parts of 3-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-3H-imidazo[4,5-b]pyridine dihydrochloride, 5.3 parts of sodium carbonate and 120 parts of 4-methyl-2-pentanone was stirred and refluxed for 15 minutes using a water separator. 3.2 Parts of 2-ethenylpyridine were added at reflux temperature and stirring was continued for 3 hours at reflux using a water separator. Then there were added 3.2 parts of 2-ethenylpyridine and the whole was stirred and refluxed for 19.50 hours using a water separator. After cooling, the salts were filtered off and the filtrate was washed with water, dried, filtered and evaporated. The residue was converted into the ethanedioate salt in 2-propanone. The salt was filtered off and crystallized from a mixture of ethanol and 2-propanone, yielding 2.5 parts (17%) of 3-[(4-fluorophenyl)methyl]-2-[[1-[2-(2-pyridinyl)ethyl]-4-piperidinyl]methyl]-3H-imidazo[4,5-b]pyridine ethanedioate (1:3); mp. 143.1°C (101).

In a similar manner there were also prepared:

4-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]-2-butanone ethanedioate(2:5).; mp. 163.4°C (102); and 1-[(4-fluorophenyl)methyl]-2-[[1-[2-(2-pyridinyl)ethyl]-4-piperidinyl]-methyl]-1H-benzimidazole ethanedioate(1:3).monohydrate; mp. 138.3°C (103).

Example 38

1.8 Parts of gaseous oxirane were bubbled through a stirred mixture of 8.5 parts of 1-[(4-fluorophenyl)methyl]-2-[(4-piperidinyl)thio]-1H-benzimidazole and 120 parts of methanol. Stirring was continued overnight at room temperature. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in 2-propanone. The salt was filtered off and dried, yielding 5.5 parts (48%) of 4-[[1-[(4-fluorophenyl)-methyl]-1H-benzimidazol-2-yl]thio]-1-piperidineethanol ethanedioate (1:1); mp. 165.2°C (104).

In a similar manner there were also prepared:

- 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]oxy]-1-piperidine-ethanol (E)-2-butenedioate (2:3); mp. 156.1°C (105);
- 4-[(1-phenyl-1H-benzimidazol-2-yl)methyl]-1-piperidineethanol;
- 5 mp. 112.2°C (106);
- 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidineethanol; mp. 135.6°C (107);
- 4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidineethanol (108);
- 10 4-[[1-(4-thiazolylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidineethanol ethanedioate (2:5); mp. 123.5-137.8°C (109);
- 4-[[1-[(4-methoxyphenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidineethanol ethanedioate (1:2); mp. 148.5°C (110);
- 4-[[3-(2-pyridinylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-
- 15 piperidineethanol (E)-2-butenedioate (2:3); mp. 151.0°C (111);
- 4-[[1-(phenylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidineethanol; mp. 136.9°C (112);
- 4-[[1-[(4-methylphenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidineethanol ethanedioate (1:2); mp. 167.7°C (113);
- 20 4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-4-[[1-(naphthalenyloxy)methyl]-1-piperidineethanol (E)-2-butenedioate (2:3); mp. 144.7°C (114);
- 4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidineethanol; mp. 116.8°C (115); and
- 25 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-4-[(2-naphthalenyloxy)methyl]-1-piperidineethanol ethanedioate (1:2); mp. 152.9°C (116).

Example 39

- A mixture of 7.9 parts of 3-[(4-fluorophenylmethyl)-2-(4-
- 30 piperidinylmethyl)-3H-imidazo[4,5-b]pyridine dihydrochloride, 8.5 parts of sodium carbonate and 120 parts of 4-methyl-2-pentanone was stirred and refluxed for 30 minutes using a water separator. 7.8
- Parts of 2-thiopheneethanol methanesulfonate (ester) were added and the whole was stirred and refluxed for 4 hours using a water
- 35 separator. After cooling, the salts were filtered off, washed with

4-methyl-2-pentanone and the filtrate was washed with water. The organic layer was dried, filtered and evaporated. The residue was converted into the hydrochloride salt in 2-propanol and 2-propanone. The salt was filtered off and dried in vacuo at 60°C, yielding 8.0 parts (76%) of 3-[(4-fluorophenyl)methyl]-2-[[1-[2-(2-thienyl)ethyl]-4-piperidinyl]methyl]-3H-imidazo[4,5-b]pyridine dihydrochloride monohydrate; mp. 210.8°C (117).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there was also prepared:

10 1-[(4-fluorophenyl)methyl]-2-[[1-[2-(2-thienyl)ethyl]-4-piperidinyl]methyl]-1H-benzimidazole ethanedioate (1:2).hemihydrate; mp. 142.0°C (118).


Example 40

A mixture of 6.5 parts of 1-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-1H-benzimidazole, 4.2 parts of sodium carbonate and 120 parts of 4-methyl-2-pentanone was stirred and refluxed for 30 minutes using a water separator. 5.2 Parts of 1-(3-chloropropoxy)-4-fluorobenzene were added at reflux temperature and stirring was continued for 3 hours at this temperature using a water separator. After cooling to room temperature, the salts were filtered off and the filtrate was washed twice with water, dried, filtered and evaporated. The residue was converted into the ethanedioate salt in 2-propanone. The salt was filtered off, washed with 2-propanone and crystallized from methanol. The product was filtered off and dried in vacuo at 80°C, yielding 7 parts (53%) of 2-[[1-[3-(4-fluorophenoxy)propyl]-4-piperidinyl]methyl]-1-[(4-fluorophenyl)methyl]-1H-benzimidazole ethanedioate (1:2); mp. 186.7°C (119).

30

35

Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:



No.	L'	s	R¹	A¹	salt/ base	mp. (°C)
10						
120	C ₆ H ₅ O-	2	H	CH	base	143-144.5
121	4-F-C ₆ H ₄ -	3	H	CH	base	140-144
122	C ₆ H ₅ -	2	H	CH	base	183-187
15		2	4-F-C ₆ H ₄ CH ₂	CH	2(COOH) ₂	127.6
124	4-morpholinyl	2	4-F-C ₆ H ₄ CH ₂	N	2(COOH) ₂	205.2
125		2	4-F-C ₆ H ₄ CH ₂	N	2(COOH) ₂	182.2
20	C ₂ H ₅ O-	2	4-F-C ₆ H ₄ CH ₂	N	2HCl H ₂ O	180.0
127	4-F-C ₆ H ₄ C(O)-	3	4-F-C ₆ H ₄ CH ₂	N	2HCl H ₂ O	167.1
128	(CH ₃) ₂ CHNHC(O)-	1	4-F-C ₆ H ₄ CH ₂	N	(COOH) ₂	227.5
25	C ₆ H ₅ S-	2	4-F-C ₆ H ₄ CH ₂	CH	2(COOH) ₂	173.5
130	C ₆ H ₅ SO ₂	2	4-F-C ₆ H ₄ CH ₂	CH	2(COOH) ₂	193.0
131	4-morpholinyl	2	4-F-C ₆ H ₄ CH ₂	CH	*	207.7
132	1H-benzimidazol-2-yl	1	4-F-C ₆ H ₄ CH ₂	N	3(COOH) ₂	165.5
133	C ₂ H ₅ O-	2	4-F-C ₆ H ₄ CH ₂	CH	***	113.0
134	(CH ₃) ₂ CHNHC(O)-	1	4-F-C ₆ H ₄ CH ₂	CH	***	151.0
30	4-F-C ₆ H ₄ O-	3	4-F-C ₆ H ₄ CH ₂	N	2(COOH) ₂	157.1
136	1H-benzimidazol-2-yl	1	4-F-C ₆ H ₄ CH ₂	CH	**	205.1
137	2,3-dihydro-1,4-benzodioxin-2-yl	1	4-F-C ₆ H ₄ CH ₂	N	2(COOH) ₂	176.8

35

* : (E)-2-butenedioate salt (1:2)

** : (E)-2-butenedioate salt (2:3)

***: ethan dioate salt (2:5)

In a similar manner there was also prepared:

5 2-[[1-[[1-(1H-benzimidazol-2-ylmethyl)-1H-benzimidazol-2-yl]methyl]-4-piperidinyl]methyl]-3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]-pyridine; mp. 247.7°C (138).

Example 41

A mixture of 3.16 parts of 1-(3-chloropropyl)-1,3-dihydro-2H-
10 benzimidazol-2-one, 4.4 parts of 1-(2-furanylmethyl)-2-(4-piperidinylmethyl)-1H-benzimidazole, 2 parts of sodium hydrogen carbonate and 80 parts of ethanol was stirred and refluxed for 32 hours. The reaction mixture was cooled and filtered over Hyflo. The filtrate was evaporated. The residue was purified by column
15 chromatography over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in methanol. The salt was filtered off and dried, yielding 4.2 parts (43%) of 1-[3-[4-[[1-(2-furanylmethyl)-1H-benz-
20 imidazol-2-yl]methyl]-1-piperidinyl]propyl]-1,3-dihydro-2H-benzimidazol-2-one ethanedioate (1:2); mp. 214.7-218.4°C (139).

In a similar manner there were also prepared:

1-[3-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]oxy]-1-piperidinyl]propyl]-1,3-dihydro-2H-benzimidazol-2-one; mp. 186.7°C
25 (140);
3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]oxy]-1-piperidinyl]ethyl]-2,4(1H,3H)quinazolinedione; mp. 190.4°C (141).
3-[(4-fluorophenyl)methyl]-2-[[1-(2-propenyl)-4-piperidinyl]methyl]-3H-imidazo[4,5-b]pyridine dihydrochloride monohydrate; mp. 166.9°C
30 (142);
4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]-methyl]-N-(1-methylethyl)-1-piperidinepropanamide; mp. 134.0°C (143);
1-[(4-fluorophenyl)methyl]-2-[[1-(2-propenyl)-4-piperidinyl]methyl]-1H-benzimidazole ethanedioate(1:2); mp. 119.0°C (144);
35 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-(1-methylet-

hyl)-1-piperidinepropanamide (E)-2-butenedioate(2:3); mp. 138.3°C (145);
 3-[(4-fluorophenyl)methyl]-2-[[1-[2-(phenylsulfonyl)ethyl]-4-piperidinyl]-
 methyl]-3H-imidazo-[4,5-b]pyridine ethanedioate(2:3); mp. 159.0°C
 (146); and

- 5 3-[(4-fluorophenyl)methyl]-2-[[1-[2-(phenylthio)ethyl]-4-piperidinyl]met-
 hyl]-3H-imidazo[4,5-b]pyridine ethanedioate(1:2); mp. 190.0°C (147).

Example 42

- A mixture of 9.3 parts of 2-iodoacetamide, 20.0 parts of
 3-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-3H-imidazo[4,5-b]-
 10 pyridine dihydrochloride, 17.0 parts of sodium hydrogen carbonate and
 200 parts of ethanol was stirred for 3 hours at at reflux temperature.
 The salts were filtered off and the filtrate was evaporated. The
 residue was purified by column chromatography over silica gel using a
 mixture of trichloromethane, methanol and ammonium hydroxide (90:9:1 by
 15 volume) as eluent. The pure fractions were collected and the eluent was
 evaporated. The residue was crystallized from a mixture of 2,2'-oxybis-
 propane and 2-propanone. The product was filtered off and dried in
 vacuo at 60°C, yielding 8.5 parts (44.5%) of 4-[[3-[(4-fluorophenyl)-
 methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidineacetamide;
 20 mp. 153.4°C (148).

In a similar manner there was also prepared:

4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-
 piperidineacetamide; mp. 187.5°C (149).

Example 43


- 25 A mixture of 5.55 parts of N-(dihydro-3,3-diphenyl-2(3H)-furanyl-
 idene)-N-methylmethanaminium bromide, 4.85 parts of 1-[(4-fluorophenyl)-
 methyl]-2-(4-piperidinylmethyl)-1H-benzimidazole, 2 parts of sodium
 carbonate and 90 parts of N,N-dimethylformamide was stirred overnight
 at 70°C. The reaction mixture was poured into water. The product was
 30 extracted with 4-methyl-2-pentanone. The extract was dried, filtered
 and evaporated. The residue was purified by column chromatography over
 silica gel using a mixture of trichloromethane and methanol, saturated
 with ammonia, (95:5 by volume) as eluent. The pure fractions were
 collected and the eluent was evaporated. The residue was crystallized


from acetonitrile. The product was filtered off and dried, yielding 1.8 parts (20%) of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N,N-dimethyl- α,α -diphenyl-1-piperidinebutanamide; mp. 151.4°C (150).









Example 44

A mixture of 6.62 parts of 6-(2-bromoethyl)-3,7-dimethyl-5H-thiazolo[3,2-a]pyrimidin-5-one monohydrobromide, 4.45 parts of 3-(2-furanylmethyl)-2-(4-piperidinylmethyl)-3H-imidazo[4,5-b]-pyridine, 4.8 parts of sodium carbonate and 90 parts of N,N-dimethylformamide was stirred and heated overnight at 70°C. The reaction mixture was poured into water. The product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the hydrochloride salt in ethanol. The salt was filtered off and dried, yielding 4.5 parts (48%) of 6-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-3,7-dimethyl-5H-thiazolo[3,2-a]pyrimidin-5-one trihydrochloride. monohydrate; mp. 248.6°C (151).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:



No.	L'	s	B	R ^{1-a}	A ¹	salt/base	mp.(°C)
152		3	CH ₂	2-furanyl-	N	1 1/2 (COOH) ₂	206.2

No.	L'	s	B	R ^{1-a}	A ¹	salt/base	mp. (°C)
5							
153		3	CH ₂	4-F-C ₆ H ₄ -	N	*	132.6
10							
154		3	CH ₂	4-CH ₃ O-C ₆ H ₄ -	CH	**	168.7
15							
155		3	CH ₂	C ₆ H ₅ -	CH	2(COOH) ₂	211.1
20							
156		3	CH ₂	C ₆ H ₅ -	N	3(COOH) ₂	147.5
25							
157		3	CH ₂	4-F-C ₆ H ₄ -	CH	base	186.6
30							
158		2	CH ₂	4-F-C ₆ H ₄ -	CH	***	192.6
35							
159		3	CH ₂	2-furanyl-	CH	base	179.1
160		2	CH ₂	4-F-C ₄ H ₆ -	CH	**	194.9

5

10








15

20








25

30

35


No.	L'	s	B	R ^{1-a}	A ¹	salt/base	mp. (°C)
161		2	CH ₂	4-F-C ₆ H ₄ -	CH	***	174.7
162		2	CH ₂	4-F-C ₆ H ₄ -	CH	**	186.9
163		2	CH ₂	2-furanyl	CH	2(COOH) ₂ .2H ₂ O	164.7
164		2	CH ₂	4-F-C ₆ H ₄ -	CH	base	168.6
165		2	CH ₂	2-furanyl-	CH	2HCl.H ₂ O	240.1
166		2	CH ₂	2-furanyl-	CH	3HCl.2H ₂ O	197.4
167		2	CH ₂	2-furanyl	CH	3HCl.2H ₂ O	215.8

0151826


No.	L'	s	B	R ^{1-a}	A ¹	salt/base	mp. (°C)
168		2	CH ₂	2-furanyl	N	3HCl.H ₂ O	250.2
169		3	CH ₂	2-furanyl	N	base	198.2
170		2	CH ₂	2-furanyl	N	2HCl.H ₂ O	227.4
171		2	CH ₂	2-furanyl	N	base	199.2
172		3	CH ₂	4-F-C ₆ H ₄ -	N	base	183.6
173		2	CH ₂	2-furanyl-	N	3HCl.2H ₂ O	186.8 (dec.)
174		2	CH ₂	2-furanyl-	N	2HCl.H ₂ O	204.3 (dec.)

5


No.	L ¹	S	B	R ^{1-a}	A ¹	salt/base	mp. (°C)
-----	----------------	---	---	------------------	----------------	-----------	----------

175		2	CH ₂	2-furanyl	CH	base	175.2
-----	---	---	-----------------	-----------	----	------	-------


10

176		2	CH ₂	2-furanyl	N	2HCl.H ₂ O	182.1
-----	---	---	-----------------	-----------	---	-----------------------	-------


15

177		2	CH ₂	4-F-C ₆ H ₄ -	N	3HCl.H ₂ O	229.7
-----	---	---	-----------------	-------------------------------------	---	-----------------------	-------


20


178		2	CH ₂	2-furanyl-	N	***	183.6
-----	--	---	-----------------	------------	---	-----	-------

25







179		2	CH ₂	4-F-C ₆ H ₄ -	N	2HCl.H ₂ O	240.9
-----	---	---	-----------------	-------------------------------------	---	-----------------------	-------

30

180		2	CH ₂	2-furanyl-	N	2 1/2 (COOH) ₂ . H ₂ O	163.1
-----	---	---	-----------------	------------	---	--	-------

181		2	CH ₂	4-F-C ₆ H ₄ -	CH	2 1/2 (COOH) ₂	161.0
-----	---	---	-----------------	-------------------------------------	----	---------------------------	-------

35


No.	L'	s	B	R ^{1-a}	A ¹	salt/base	mp. (°C)
182		2	CH ₂	4-F-C ₆ H ₄ -	CH	base	101.2
183		1	CH ₂	4-F-C ₆ H ₄ -	CH	base	164.3
184		1	CH ₂	4-F-C ₆ H ₄ -	CH	3(COOH) ₂	161.4
185		2	O	4-F-C ₆ H ₄ -	CH	2HCl	194.8
186		2	CH ₂	2-thienyl-	CH	base	196.0
187		2	CH ₂	4-thiazolyl-	CH	base	210.6

* : (E)-2-butenedioate salt (2:3).monohydrate

** : (E)-2-butenedioate salt (2:3)

*** : (E)-2-butenedioate salt (1:2)

In a similar manner there were also prepared:



Comp. No.	L'	s	B	R ^{1-a}	A ¹	base or salt	mp. °C
188	(4-F-C ₆ H ₄) ₂ -CH-	3	O	4-F-C ₆ H ₄ -	CH	2HNO ₃	140.3
189	4-CH ₃ O-C ₆ H ₄ -	2	CH ₂	2-furanyl-	N	2(COOH) ₂	159.4
190	4-CH ₃ O-C ₆ H ₄ -	2	O	4-F-C ₆ H ₄ -	CH	*	155.2
191	4-CH ₃ O-C ₆ H ₄ -	2	CH ₂	4-CH ₃ OC ₆ H ₄ -	CH	**	223.8
192	(4-F-C ₆ H ₄) ₂ -CH-	3	CH ₂	2-furanyl-	N	2(COOH) ₂	144.1
193	(4-F-C ₆ H ₄) ₂ -CH-	3	CH ₂	2-pyridinyl-	N	2(COOH) ₂	134.9
194	(4-F-C ₆ H ₄) ₂ -CH-	3	CH ₂	4-CH ₃ OC ₆ H ₄ -	CH	2(COOH) ₂	126.7
195	(4-F-C ₆ H ₄) ₂ -CH-	3	CH ₂	4-F-C ₆ H ₄ -	N	2(COOH) ₂	182.2
196	(4-F-C ₆ H ₄) ₂ -CH-	3	CH ₂	C ₆ H ₅ -	CH	2 1/2(COOH) ₂	140.4
197	(4-F-C ₆ H ₄) ₂ -CH-	3	CH ₂	C ₆ H ₅ -	N	2(COOH) ₂	190.2
198	4-CH ₃ O-C ₆ H ₄ -	2	CH ₂	C ₆ H ₅ -	N	*.H ₂ O	123.4
199	2,3-dihydro-1,4-benzodioxin-2-yl	1	CH ₂	C ₆ H ₅ -	CH	2 1/2(COOH) ₂	224.5
200	4-CH ₃ O-C ₆ H ₄ -	2	S	4-F-C ₆ H ₄ -	CH	2 1/2(COOH) ₂	148.1
201	C ₂ H ₅ O-C(=O)-	1	CH ₂	4-F-C ₆ H ₄ -	CH	2HCl.H ₂ O	174.7

* : (E)-2-butenedioate(1:2)

** : (E)-2-butenedioate(1:1)

0 In a similar manner there were also prepared:

1-[(4-fluorophenyl)methyl]-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-methyl]-1H-imidazo-[4,5-c]pyridine ethanedioate(1:2);
mp. 193.0°C (202).

5 1-[(4-fluorophenyl)methyl]-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-

piperidinyl)methyl]-1H-imidazo-[4,5-b]pyridine ethanedioate(1:1);
mp. 176.7°C (203).

3-[(4-fluorophenyl)methyl]-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-
piperidinyl)methyl]-3H-imidazo-[4,5-c]pyridine ethanedioate(1:2);
5 mp. 191.6°C (204).

Example 45


A mixture of 3.14 parts of 7-(2-bromoethyl)-3,4-dihydro-8-methyl-
2H,6H-pyrimido[2,1-b][1,3]thiazin-6-one monohydrobromide, 3.5 parts
of 3-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-3H-imidazo-
10 [4,5-b]pyridin, 4 parts of sodium carbonate, 0.1 parts of potassium
iodide and 90 parts of N,N-dimethylformamide was stirred and heated
overnight at 70°C. After cooling, water was added. The product was
extracted with 4-methyl-2-pentanone. The extract was dried, filtered
and evaporated. The residue was purified by column chromatography
15 over silica gel using a mixture of trichloromethane and methanol,
saturated with ammonia, (95:5 by volume) as eluent. The pure
fractions were collected and the eluent was evaporated. The residue
was converted into the ethanedioate salt in ethanol. The salt was
filtered off and dried, yielding 3.7 parts (61%) of 7-[2-[4-[[3-[(4-
20 fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1-piperi-
dinyl]ethyl]-3,4-dihydro-8-methyl-2H,6H-pyrimido[2,1-b][1,3]thiazin-
6-one ethanedioate (1:2); mp. 190.5°C (205).

Following the same procedure and using equivalent amounts of the
appropriate starting materials, there were also prepared:

25

30

35



5

10








15









20







25

30

35

No.	L'	s	B	R ¹	A ¹	salt/base	mp. (°C)
206		3	CH ₂	C ₆ H ₅ -	CH	base	160.7
207		3	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH	**	145.7
208		2	CH ₂	2-furanyl-methyl-	CH	base	210.7
209		2	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH	base	177.8
210		1	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH	2(COOH) ₂	198.6- 200.1
211		3	CH ₂	4-CH ₃ -C ₆ H ₄ CH ₂ -	CH	2(COOH) ₂ ·H ₂ O	166.2
212		1	CH ₂	2-thienyl-methyl-	N	(COOH) ₂	184.1

No.	L'	s	B	R ¹	A ¹	salt/base	mp. (°C)
213		1	CH ₂	4-CH ₃ -C ₆ H ₄ -CH ₂ -	CH	* .1/2H ₂ O	202.0
214		2	CH ₂	4-F-C ₆ H ₄ -CH ₂ -	CH	2(COOH) ₂	195.8
215		3	CH ₂	2-thienyl-methyl-	N	1 1/2(COOH) ₂	173.3
216		2	CH ₂	2-furanyl-methyl-	CH	base	155.5
217		1	CH ₂	2-pyridinyl-methyl-	N	2 1/2(COOH) ₂	157.2
218		2	CH ₂	2-furanyl-methyl-	CH	2 1/2(COOH) ₂	115.2
219		2	CH ₂	4-F-C ₆ H ₄ -CH ₂ -	CH	2(COOH) ₂	139.8
220		2	CH ₂	4-F-C ₆ H ₄ -CH ₂ -	CH	base	173.8


No.	L ¹	s	B	R ¹	A ¹	salt/base	mp. (°C)
5							
221		2	CH ₂	2-furanyl-methyl-	CH	2(COOH) ₂ ·H ₂ O	162.8
10							
222		2	CH ₂	2-furanyl-methyl-	CH	*	192.4
15							
223		2	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH	*	212.1
20							
224		2	CH ₂	4-F-C ₆ H ₄ CH ₂ -	N	base	192.7
25							
225		2	CH ₂	4-F-C ₆ H ₄ CH ₂ -	N	2 1/2(COOH) ₂ ·H ₂ O	125.6
30							
226		2	CH ₂	2-furanyl-methyl	CH	3(COOH) ₂ ·H ₂ O	125.6

30 * : (E)-2-butenedioate (2:3)

** : (E)-2-butenedioate (1:2)

35

In a similar manner there were also prepared:



5

Comp. No.	L'	s	R¹	A¹	base or salt	mp. °C
10227	4-CH₃O-C₆H₄-	2	4-F-C₆H₄CH₂-	CH	*	190.5
228	4-F-C₆H₄-CO-	3	4-F-C₆H₄CH₂-	N	*	152.6
229	4-CH₃O-C₆H₄-	2	2-furanylmethyl	CH	***	209.0
230	C₆H₅-CH=CH-	2	2-furanylmethyl	CH	**	167.8
231	4-CH₃O-C₆H₄-	2	C₆H₅-	CH	*	195.8
15232	4-CH₃O-C₆H₄-	2	2-thienylmethyl	CH	***	205.9- 207.6
233	(4-F-C₆H₄)₂CH-	3	C₆H₅-	CH	2(COOH)₂	163.8- 165.6
234	4-CH₃O-C₆H₄-	2	2-pyridinylmethyl	N	**	170.5
20235	(4-F-C₆H₄)₂CH-	3	4-F-C₆H₄CH₂-	CH	2(COOH)₂	130.1
236	4-CH₃O-C₆H₄-	2	4-F-C₆H₄CH₂-	N	2(COOH)₂	155.7
237	C₂H₅-O-CO-NH-	2	2-thienylmethyl	N	base	-
238	4-CH₃O-C₆H₄-	2	2-thienylmethyl	N	***	198.0
239	4-CH₃O-C₆H₄-	2	4-CH₃C₆H₄CH₂	CH	***	214.5
25240	(4-F-C₆H₄)₂CH-	3	2-thienyl₁₄CH₂-	N	(COOH)₂	192.7
241	(4-F-C₆H₄)₂CH-	3	4-CH₃C₆H₄CH₂	CH	2 1/2(COOH)₂	116.1
242	4-CH₃O-C₆H₄-	2	CH₃	CH	2(COOH)₂	164.3

* : (E)-2-butenedioate (2:3)

30 ** : (E)-2-butenedioate (1:2)

***: (E)-2-butenedioate (1:1)

In a similar manner there were also prepared:

ethyl [2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]-thio]-1-piperidinyl)ethyl]carbamate dihydrobromide.hemihydrate; mp.


191.4°C (243); and

3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]thio]-1-piperidinyl]-ethyl]-2-methyl-4H-pyrido[1,2-a]pyrimidin-4-one trihydrochloride.monohydrate; mp. 177.8°C (244).

5 Example 46

A mixture of 1.9 parts of 1-(2-chloroethyl)-4-methoxybenzene, 4 parts of 1-(4-piperidinylmethyl)-1-(4-thiazolylmethyl)-1H-benzimidazole trihydrobromide, 8 parts of sodium carbonate, 0.1 parts of sodium iodide and 45 parts of N,N-dimethylacetamide was stirred overnight at 75°C. After cooling, the reaction mixture was poured into water and the product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residual oil was converted into the hydrochloride salt in 2-propanol and the whole was evaporated. The sticky residue was crystallized from acetonitrile, yielding 2.54 parts of 2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1-(4-thiazolylmethyl)-1H-benzimidazole dihydrochloride.monohydrate; mp. 154.6-156.1°C (245).







In a similar manner there were also prepared:








25

No.	L'	s	B	R¹-a	A¹	salt/base	mp.(°C)
246	(CH₃)₂CH-	0	CH₂	4-thiazolyl- methyl-	CH	2(COOH)₂	187.5-189.7
247	(4-F-C₆H₄)₂-CH-	3	CH₂	4-thiazolyl- methyl-	CH	2(COOH)₂	193.3-195.3
248	C₂H₅O(C=O)NH-	2	CH₂	4-thiazolyl- methyl-	CH	2(COOH)₂.H₂O	181.1


35

No.	L'	s	B	R ¹ -a	A ¹	salt/base	mp. (°C)
249		3	CH ₂	2-pyridinyl- N methyl-		2(COOH) ₂ .H ₂ O	159.6
250	C ₆ H ₅ CH=CH-	1	CH ₂	H	CH	base	199.2
251	4-CH ₃ O-C ₆ H ₄ -	2	CH ₂	H	CH	base	-
252	(CH ₃) ₂ CH-NH-C(=O)-	1	CH ₂	H	CH	base	201.3
253	C ₂ H ₅ OC(=O)-	1	CH ₂	4-F-C ₆ H ₄ CH ₂ - N.		2HCl.H ₂ O	147.8
254	4-CH ₃ O-C ₆ H ₄ -	2	SO ₂	4-F-C ₆ H ₄ CH ₂ - CH		base	111.3
255		3	S	4-F-C ₆ H ₄ CH ₂ - CH		base	140.6
256		2	CH ₂	2-pyridinyl- N methyl-		base	151.6
257		2	CH ₂	2-thienyl- methyl-	CH	base	127.0
258		2	CH ₂	2-pyridinyl- N methyl-		base	178.7
259		2	CH ₂	2-thienyl- methyl-	CH	2(COOH) ₂	216.1

No.	L'	s	B	R ¹ -a	A ¹	salt/base	mp. (°C)
260	4-CH ₃ O-C ₆ H ₄ -	2	CH ₂	3-furanyl-methyl-	CH	2 1/2(COOH) ₂	161.3
261		2	CH ₂	4-thiazolyl-methyl-	CH	base	179.5
262		2	CH ₂	4-thiazolyl-methyl-	CH	2 1/2(COOH) ₂ H ₂ O	187.2
263		2	CH ₂	2-thienyl-methyl-	CH	2(COOH) ₂	194.3
264		2	O	4-F-C ₆ H ₄ CH ₂ -	CH	*	149.8
265		2	CH ₂	4-thiazolyl-methyl-	CH	base	138.2
266	4-CH ₃ O-C ₆ H ₄ -	2	CH ₂	5-methyl-2-furanylmethyl-	CH	2(COOH) ₂	179.9

*: (E)-2-butenedioate salt(1:2)

In a similar manner there were also prepared:



5




10

15

20

25

30

No.	L'	s	R	B	A ₁ =A ₂ -A ₃ =A ₄	iso- meric form	salt/base	mp. °C
267	4-CH ₃ O-C ₆ H ₄ -	2	CH ₃	O	CH=CH-CH=CH	cis	2(COOH) ₂	160.2
268	4-CH ₃ O-C ₆ H ₄ -	2	CH ₃	O	-CH=CH-CH=CH	trans	base	135.6
269	4-CH ₃ O-C ₆ H ₄ -	2	H	CH ₂	CH=CH-C=CH CH ₃	-	**	192.7
270		2	H	CH ₂	CH=N-CH=CH	-	3(COOH) ₂	155.7
271	4-CH ₃ O-C ₆ H ₄ -	2	H	CH ₂	CH=CH-C=CH OCH ₃	-	2HCl. 1 1/2H ₂ O	192.7
272		2	H	CH ₂	CH=CH-CH=N	-	2(COOH) ₂ 1/2H ₂ O	155.7
273	4-CH ₃ O-C ₆ H ₄ -	2	H	CH ₂	CH=C-CH=CH OCH ₃	-	2(COOH) ₂	163.9
274		2	H	CH ₂	CH=CH-N=CH	-	2(COOH) ₂ H ₂ O	178.3
275	4-CH ₃ O-C ₆ H ₄ -	2	H	CH ₂	CH=CH-C=CH OH	-	***	215.5
276	4-CH ₃ O-C ₆ H ₄ -	2	H	CH ₂	CH=C-CH=CH OH	-	base	210.7

** : (E)-2-butenedioate salt (2:3)

***: (E)-2-butenedioate salt (1:1)

35

Example 47

A mixture of 13.3 parts of 1-(2-chloroethyl)-4-methoxybenzene, 23.8 parts of 5(or 6)-fluoro-1-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-1H-benzimidazole, 14.8 parts of sodium carbonate, 5 0.5 parts of potassium iodide and 250 parts of N,N-dimethylacetamide was stirred at 100°C for 5 hours. After cooling, the mixture was poured into ice water. This mixture was extracted three times with methylbenzene. The combined organic layers were washed twice with water, dried, filtered and evaporated. The residue was purified by 10 column chromatography over silica gel using a mixture of ethyl acetate, ethanol and ammonia (96:4:1 by volume) as eluent. The first fraction was collected and the eluent was evaporated. The residue was converted into the hydrochloride salt in 2,2'-oxybispropane and 2-propanol. The salt was filtered off and crystallized from a 15 mixture of 2-propanol and 2,2'-oxybispropane, yielding 9.2 parts (48%) of 5-fluoro-1-[(4-fluorophenyl)methyl]-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl)methyl]-1H-benzimidazole dihydrochloride dihydrate; mp. 101.9°C (277).

The second fraction was collected and the eluent was evaporated. The 20 residue was converted into the (E)-2-butenedioate salt in 2,2'-oxybispropane and 2-propanol. The salt was filtered off and crystallized from a mixture of 2-propanol and 2,2'-oxybispropane, yielding 6 parts (25%) of 6-fluoro-1-[(4-fluorophenyl)methyl]-2-(4-piperidinylmethyl)-1H-benzimidazole; mp. 191.5°C (278).

25 Example 48

A mixture of 6.4 parts of 2-chloroacetonitrile, 27 parts of 3-[(4-methylphenyl)methyl]-2-(4-piperidinylmethyl)-3H-imidazo[4,5-b]-pyridine, 13 parts of sodium carbonate and 450 parts of N,N-dimethylformamide was stirred overnight at room temperature. The reaction mixture was poured into water. The product was extracted 30 with 4-methyl-2-pentanone. The extract was washed with water, dried, filtered and evaporated. The residue was crystallized from 1,1'-oxybisethane, yielding 19 parts (62%) of 4-[[3-[(4-methylphenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1-piperidineacetonitrile; mp. 131.3°C (279).

Following the same procedure and using equivalent amounts of

the appropriate starting materials; there were also prepared:


4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinebutanenitrile (280);

4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidine-
5 butanenitrile (281);

4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinebutanenitrile (282).

In a similar manner there were also prepared:

10



15

No.	R	B	R ₁	A ₁ =A ₂ -A ₃ =A ₄	mp. °C
283	H	CH ₂	4-F-C ₆ H ₄ -CH ₂	CH=CH-CH=CH	146.1
284	H	CH ₂	C ₆ H ₅	CH=CH-CH=CH	141.4
285	H	CH ₂	2-furanylmethyl	CH=CH-CH=CH	152.5
20 286	H	O	4-F-C ₆ H ₄ -CH ₂	CH=CH-CH=CH	-
287	H	CH ₂	2-thienylmethyl	CH=CH-CH=CH	-
288	H	CH ₂	3-pyridinylmethyl	CH=CH-CH=CH	-
289	H	CH ₂	4-thiazolylmethyl	CH=CH-CH=CH	91.2-93.0
290	H	CH ₂	4-F-C ₆ H ₄ -CH ₂	N=CH-CH=CH	98.9
25 291	H	CH ₂	2-furanylmethyl	N=CH-CH=CH	124.2
292	H	CH ₂	2-pyridinylmethyl	N=CH-CH=CH	137.9
293	H	CH ₂	4-CH ₃ O-C ₆ H ₄ -CH ₂	CH=CH-CH=CH	129.8
294	H	CH ₂	H	CH=CH-CH=CH	205.4
295	H	CH ₂	4-CH ₃ -C ₆ H ₄ -CH ₂	CH=CH-CH=CH	161.6
30 296	H	CH ₂	C ₆ H ₅ -CH ₂	N=CH-CH=CH	140.0
297	H	CH ₂	C ₆ H ₅ -CH ₂	CH=CH-CH=CH	174.3
298	H	CH ₂	4-CH ₃ O-C ₆ H ₄ -CH ₂	N=CH-CH=CH	96.6
299	H	CH ₂	4-F-C ₆ H ₄ -CH ₂	CH=CH-CH=N	-
300*	CH ₃	CH ₂	4-F-C ₆ H ₄ -CH ₂	CH=CH-CH=CH	127.4
35 301	H	CH ₂	4-F-C ₆ H ₄ -CH ₂	CH=N-CH=CH	132.9

* : cis-isomer

Example 49

A mixture of 7.4 parts of 4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinebutanenitrile and 240 parts of methanol saturated with ammonia was hydrogenated at normal pressure and at room temperature with 3 parts of Raney-nickel catalyst. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated, yielding 7.33 parts (99%) of 4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinebutanamine (302).


Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:

- 4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidineethanamine (E)-2-butenedioate (1:3); mp. 210.9°C (303);
- 4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidineethanamine (E)-2-butenedioate (1:3); mp. 203.0°C (304);
- 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinebutanamine (305);
- 4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinebutanamine (306);
- 4-[[1-[(4-fluorophenyl)methyl]-1H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidineethanamine (307);
- 4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-c]pyridin-2-yl]methyl]-1-piperidineethanamine (308);
- cis-4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]oxy]-3-methyl-1-piperidineethanamine (309).

30

35

In a similar manner there were also prepared:



Comp. No.	B	R ¹	A ¹
10 310	CH ₂	4-F-C ₆ H ₄ -CH ₂ -	CH
311	CH ₂	C ₆ H ₅ -	CH
312	O	4-F-C ₆ H ₄ -CH ₂ -	CH
313	CH ₂	2-thienylmethyl	CH
314	CH ₂	3-pyridinylmethyl	CH
15 315	CH ₂	4-thiazolylmethyl	CH
316	CH ₂	4-F-C ₆ H ₄ -CH ₂ -	N
317	CH ₂	2-pyridinylmethyl	N
318	CH ₂	4-CH ₃ O-C ₆ H ₄ -CH ₂ -	CH
319	CH ₂	H	CH
20 320	CH ₂	4-CH ₃ -C ₆ H ₄ -CH ₂ -	CH
321	CH ₂	C ₆ H ₅ -CH ₂ -	CH
322	CH ₂	C ₆ H ₅ -CH ₂ -	N
323	CH ₂	4-CH ₃ -C ₆ H ₄ -CH ₂ -	N
324	CH ₂	4-CH ₃ O-C ₆ H ₄ -CH ₂ -	N
25			

Example 50

A mixture of 20.7 parts of ethyl [2-[4-[[3-(2-thienylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]carbamate, 22.1 parts of potassium hydroxide and 200 parts of 2-propanol was stirred and refluxed overnight. The reaction mixture was evaporated. Water was added to the residue. The product was extracted with dichloromethane. The extract was dried, filtered and evaporated, yielding 13 parts (76%) of 4-[[[3-(2-thienylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinethanamine (325).

In a similar manner there was also prepared:

4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]thio]-1-piperidineethanamine (326).

Example 51

- 5 A mixture of 12 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1'-(phenylmethyl)[1,4'-bipiperidine] and 200 parts of methanol was hydrogenated at normal pressure and 50°C with 3 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and
10 the filtrate was evaporated. The residue was converted into the (E)-2-butenedioate salt in methanol. The salt was filtered off and dried, yielding 7.87 parts (51.3%) of 4-[[1-[(4-fluorophenyl)-methyl]-1H-benzimidazol-2-yl]methyl]-[1,4'-bipiperidine] (E)-2-butenedioate(1:2); mp. 226.9°C (327).

15 Example 52

- A mixture of 3 parts of 2-chloro-1H-benzimidazole, 7.3 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidineethanamine and 0.1 parts of potassium iodide was stirred for 1 hour in an oil bath at 160°C. After cooling to room temperature, the
20 whole was pulverized and stirred in a mixture of water, ammonium hydroxide and trichloromethane. The layers were separated. The organic layer was washed twice with water, dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of hexane, trichloromethane, methanol and
25 ammonium hydroxide (45:45:9:1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt in 2-propanone. The salt was filtered off and crystallized from ethanol. The product was filtered off and dried in vacuo at 70°C overnight, yielding 5 parts
30 (38.0%) of N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]1H-benzimidazol-2-amine (E)-2-butenedioate(2:3); mp. 134.5°C (328).

In a similar manner there were also prepared:

- N-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]-methyl]-1-piperidinyl]ethyl]1H-benzimidazol-2-amine; mp. 161.9°C (329).
35

Example 53

A mixture of 3 parts of 2-methylthio-thiazolo[5,4-b]pyridine and 5.5 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]-methyl]-1-piperidineethanamine was stirred for 24 hours at 140°C.

- 5 The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in ethanol. The salt was filtered off and dried, yielding 1 part (8.6%) of N-[2-[4-[[1-[(4-fluorophenyl)-methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]thiazolo[5,4-b]pyridin-2-amine ethanedioate(1:3); mp. 148.2°C (330).


In a similar manner there were also prepared:

- 2-[[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]amino]-4(1H)-pyrimidinone; mp. 164.0°C (331);
15 N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]thiazolo[4,5-c]pyridin-2-amine ethanedioate(1:3); mp. 188.0°C (332).

Example 54

- 20 A mixture of 1.7 parts of 2-chloropyrimidine, 5.7 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]thio]-1-piperidineethanamine, 1.3 parts of sodium hydrogen carbonate and 120 parts of ethanol was stirred and refluxed overnight. The reaction mixture was filtered and the filtrate was evaporated. The residue
25 was taken up in trichloromethane. The solution was washed with water, dried, filtered and evaporated. The residue was purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was
30 converted into the ethanedioate salt in ethanol. The salt was filtered off and dried, yielding 3.7 parts (44.6%) of N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]thio]-1-piperidinyl]ethyl]-2-pyrimidinamine ethanedioate (1:1); mp. 189.2°C (333).

In a similar manner there were also prepared:



Comp. No.	L'	B	R ^{1-a}	A ¹	base or salt	mp. °C
10 334	2-pyrimidinyl	CH ₂	2-furanyl-	CH	base	103.0
335	2-thiazolyl	CH ₂	2-furanyl-	CH	*	159.6
336	2-pyrimidinyl	CH ₂	2-thienyl-	CH	**	184.6-188.6
337	2-pyrimidinyl	CH ₂	3-pyridinyl-	CH	4(COOH) ₂	176.1-180.5
338	2-pyrimidinyl	CH ₂	4-thiazolyl-	CH	4(COOH) ₂	192.3-194.0
15 339	2-pyrimidinyl	CH ₂	2-furanyl-	N	3(COOH) ₂	107.3
340	2-pyrimidinyl	CH ₂	2-pyridinyl-	N	3 1/2 (COOH) ₂	151.7
341	2-pyrimidinyl	CH ₂	4-CH ₃ OC ₆ H ₄ -	CH	**	182.2
342	2-pyrimidinyl	CH ₂	2-thienyl-	N	**·H ₂ O	152.9
343	2-pyrimidinyl	CH ₂	4-CH ₃ -C ₆ H ₄ -	CH	2 1/2(COOH) ₂	160.5
20 344	2-pyrimidinyl	CH ₂	C ₆ H ₅ -	CH	**	194.8
345	2-pyrimidinyl	CH ₂	C ₆ H ₅ -	N	3(COOH) ₂	172.5
346	2-pyrimidinyl	O	4-F-C ₆ H ₄ -	CH	base	155.1

* : (E)-2-butenedioate salt (1:3)

25 ** : (E)-2-butenedioate salt (2:3)

In a similar manner there were also prepared:

4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1'-(2-pyrimidinyl)-[1,4'-bipiperidine] ethanedioate(2:7); mp. 169.7°C (347);

30 N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-imidazo[4,5-b]pyridin-2-yl]-methyl]-1-piperidinyl]ethyl]-2-pyrimidinamine ethanedioate(2:5); mp. 173.4°C (348);

cis-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]oxy]-3-methyl-1-piperidinyl]ethyl]-2-pyrimidinamine; mp. 94.0°C (349);

35 6-chloro-N⁴-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-

yl)methyl]-1-piperidinyl]ethyl]-4,5-pyrimidinediamine (350); and
 N-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-c]pyridin-2-yl]-
 methyl]-1-piperidinyl]-ethyl]-2-pyrimidinamine ethanedioate(1:2);
 mp. 192.5°C (351).


5 Example 55

A mixture of 3.3 parts of 2-bromothiazole, 5.09 parts of
 4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1-
 piperidineethanamine, 3 parts of sodium carbonate and 45 parts of
 N,N-dimethylacetamide was stirred overnight at 130°C. The reaction
 10 mixture was poured into water and the product was extracted twice
 with 4-methyl-2-pentanone. The extract was dried, filtered and
 evaporated. The residue was purified by column chromatography over
 silica gel using a mixture of trichloromethane and methanol
 saturated with ammonia (95:5 by volume) as eluent. The pure
 15 fractions were collected and the eluent was evaporated. The residue
 was converted into the ethanedioate salt in ethanol. The salt was
 filtered off and dried, yielding 2.5 parts (27%) of 4-[[3-(2-
 furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-N-(2-thiazolyl)-1-
 piperidineethanamine ethanedioate (1:3); mp. 173.0°C (352).

20

In a similar manner there were also prepared:

25



30

Comp. No.	L'	B	R ¹	A ¹	base or salt	mp. °C
353	2-thiazolyl	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH	base	115.6
354	2-thiazolyl	CH ₂	C ₆ H ₅ -	CH	base	59.5
355	2-thiazolyl	O	4-F-C ₆ H ₄ CH ₂ -	CH	base	132.1
356	2-thiazolyl	CH ₂	2-thienylmethyl	CH	3(COOH) ₂ H ₂ O	135.6
357	2-thiazolyl	CH ₂	4-F-C ₆ H ₄ CH ₂ -	N	2 1/2(COOH) ₂ H ₂ O	150.9

Comp. No.	L'	B	R ¹	A ¹	base or salt	mp. °C
5	358	2-pyrimidinyl	CH ₂	H	CH base	199.4
	359	2-thiazolyl	CH ₂	H	CH base	180.4
	360	5-chloro-2-pyridinyl	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH base	-
	361	2-thiazolyl	CH ₂	2-thienylmethyl-	N 3(COOH) ₂	129.4
10	362	2-thiazolyl	CH ₂	C ₆ H ₅ CH ₂ -	N 3 1/2(COOH) ₂	129.4
	363	2-thiazolyl	CH ₂	C ₆ H ₅ CH ₂ -	CH 4(COOH) ₂	142.7
	364	2-thiazolyl	CH ₂	4-CH ₃ -C ₆ H ₄ CH ₂ -	CH 3(COOH) ₂ H ₂ O	138.6
	365	2-thiazolyl	CH ₂	4-CH ₃ O-C ₆ H ₄ CH ₂ -	N base	144.7
	366	2-NO ₂ -C ₆ H ₄	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH base	120.5
15	367	2-benzothiazolyl	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH 3 1/2(COOH) ₂ 1/2 H ₂ O	167.1
	368	2-pyrazinyl	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH 2(COOH) ₂	173.6
	369	9-methyl-9H-purin-6-yl	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH 4HCl.H ₂ O	210.3
20	370	9H-purin-6-yl	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH base	186.5
	371	8-OH, 9-CH ₃ -9H-purin-6-yl	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH 2 1/2(COOH) ₂	179.7
	372	8,9-(CH ₃) ₂ -9H-purin-6-yl	CH ₂	4-F-C ₆ H ₄ CH ₂ -	CH base	134.3
25						

In a similar manner there was also prepared:

N-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-c]pyridin-2-yl]-methyl]-1-piperidinyl]ethyl]-2-pyrazinamine ethanedioate(1:2); mp.

30 157.4°C (373).

Example 56

A mixture of 1.7 parts of 2-chloropyrimidine, 5.5 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidineethanamine, 2.12 parts of sodium carbonate, 0.1 parts of
35 potassium iodide and 90 parts of N,N-dimethylformamide was stirred

overnight at 60°-70°C. Water was added and the product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) saturated with ammonia, as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from 1,1'-oxybisethane. The product was filtered off and dried, yielding 2.6 parts (40%) of N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-pyrimidinamine; mp. 125.1°C (374).

In a similar manner there were also prepared:

N-[2-[4-[[1-phenyl-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-pyrimidinamine (E)-2-butenedioate (1:1); mp. 211.4°C (375);
N-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2-pyrimidinamine (E)-2-butenedioate (2:3); mp. 162.4°C (376);
 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-(1-methyl-4-nitro-1H-imidazol-5-yl)-1-piperidineethanamine; mp. 131.1°C (377);
 6-chloro-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-3-pyridazinamine; mp. 175.5°C (378);
N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-4-nitro-3-pyridinamine, 1-oxide; mp. 135.1°C (379);
 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1'-(1-methyl-4-nitro-1H-imidazol-5-yl)-[1,4'-bipiperidine]; mp. 144.0°C (380); and
 4-chloro-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1-phthalazinamine; mp. 169.7°C (381).

30 Example 57

To a stirred and cooled (0°-10°C) mixture of 18.3 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidineethanamine, 7.5 parts of N,N-diethylethanamine and 225 parts of tetrahydrofuran was added dropwise a solution of 8.15 parts of
 35 2,6-dichloro-4-methylpyrimidine in a small amount of tetrahydro-

furan. Upon completion, stirring was continued overnight at room temperature. The reaction mixture was evaporated. Water was added to the residue and the product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was
5 purified by HPLC over silica gel using a mixture of methylbenzene, ethanol and methanol, saturated with ammonia, (85:14:1 by volume) as eluent. The first fraction was collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in ethanol. The salt was filtered off and dried, yielding 2.2 parts
10 (6.5%) of 4-chloro-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-6-methyl-2-pyrimidinamine ethanedioate (1:2); mp. 165.8°C (382).

In a similar manner there were also prepared:

2-chloro-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]-
15 methyl]-1-piperidinyl]ethyl]-6-methyl-4-pyrimidinamine; mp. 142.9°C (383);

6-chloro-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]-methyl]-1-piperidinyl]ethyl]-4-pyrimidinamine ethanedioate(2:5); mp. 174.4°C (384).

20 Example 58

A mixture of 4 parts of 5-(2-bromoethoxy)-1-methyl-1H-tetrazole, 5.5 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]-methyl]-1-piperidineethanamine, 2.3 parts of sodium carbonate and 45 parts of N,N-dimethylformamide was stirred overnight at 70°C. The
25 reaction mixture was poured into water. The product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent
30 was evaporated. The residue was converted into the ethanedioate salt in ethanol. The salt was filtered off and dried, yielding 1.3 parts (13%) of 3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]-methyl]-1-piperidinyl]ethyl]2-oxazolidinone ethanedioate(2:5); mp. 147.9°C (385).

35

Example 59

To a stirred mixture of 22 parts of 4-[[1-[(4-fluorophenyl)-methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidineethanol dihydrochloride and 450 parts of trichloromethane were added dropwise 12 parts of thionyl chloride. Upon completion, stirring was continued overnight at reflux temperature. The reaction mixture was evaporated. The residue was stirred in methylbenzene. The product was filtered off and dried, yielding 13 parts (56.6%) of 2-[[1-(2-chloroethyl)-4-piperidinyl]methyl]-1-[(4-fluorophenyl)methyl]-1H-benzimidazole dihydrochloride (386).

Example 60

A mixture of 2.0 parts of thiazolo[5,4-b]pyridine-2-thiol, 2 parts of a sodium hydride dispersion 50% and 45 parts of N,N-dimethylformamide was stirred for 2 hours. 6.5 Parts of 2-[[1-(2-chloroethyl)-4-piperidinyl]methyl]-1-[(4-fluorophenyl)methyl]-1H-benzimidazole dihydrochloride were added portionwise. Upon completion, stirring was continued over weekend. Water was added. The product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in methanol. The salt was filtered off and crystallized from methanol. The product was filtered off and dried, yielding 1.7 parts (20%) of 2-[[2-[4-[[1-[(4-fluorophenyl)-methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]thio]-thiazolo[5,4-b]pyridine ethanedioate(1:2); mp. 199.0°C (387).

In a similar manner there were also prepared:

2-[[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]thio]thiazolo[4,5-c]pyridine; mp. 121.0°C (388).

Example 61

To a stirred and warm mixture of 3.8 parts of 4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidineacetamide and 24 parts of N,N-dimethylacetamide were added portionwise 0.4 parts of a sodium hydride dispersion 59.7% at 40°C. The mixture

was heated to 80°C and stirred for 15 minutes at 80°C. 1.2 Parts of 2-chloropyrimidine were added and stirring was continued for 30 minutes at 80°C. After cooling to 40°, another 0.4 parts of a sodium hydride dispersion 59.7% were added and after stirring for 15 minutes at 80°C, another 1.2 parts of 2-chloropyrimidine were added. The whole was stirred for 30 minutes at 80°C and then cooled to 40°C. Another 0.4 parts of a sodium hydride dispersion 59.7% were added and after stirring for 15 minutes at 80°C, another 1.2 parts of 2-chloropyrimidine were added. Stirring was continued for 15 minutes at 80°C. The reaction mixture was cooled to room temperature and poured into 150 parts of water. The product was extracted three times with dichloromethane. The combined extracts were washed three times with water, dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane, methanol and methanol, saturated with ammonia, (96:3:1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt in 2-propanone. The salt was allowed to crystallize at -20°C. It was filtered off and dried, yielding 0.4 parts of 4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-N-(2-pyrimidinyl)-1-piperidineacetamide (E)-2-butenedioate(1:2); mp. 159.0°C (389).


In a similar manner there were also prepared:

1-[(4-fluorophenyl)methyl]-2-[[1-[2-(2-pyrimidinylloxy)ethyl]-4-piperidinyl]oxy]-1H-benzimidazole (E)-2-butenedioate (1:2); mp. 162.7°C (390);
 1-[(4-fluorophenyl)methyl]-2-[[1-[2-(2-pyrimidinylloxy)ethyl]-4-piperidinyl)methyl]-1H-benzimidazole ethanedioate(1:3); mp. 161.4°C (391);
 1-(2-furanylmethyl)-2-[[1-[2-(2-pyrimidinylloxy)ethyl]-4-piperidinyl]-methyl]-1H-benzimidazole ethanedioate (1:2). monohydrate; mp. 179.3°C (392);
 1-[(4-fluorophenyl)methyl]-2-[[1-[2-(2-pyrimidinylloxy)ethyl]-4-piperidinyl]thio]-1H-benzimidazole ethanedioate (1:1); mp. 186.9°C (393); and

3-[[4-fluorophenyl)methyl]-2-[[1-[2-(2-pyridinylmethoxy)ethyl]-4-piperidinyl]-methyl]-3H-imidazo[4,5-b]pyridine trihydrochloride; mp. 129.8°C (394).

Example 62

- 5 To a stirred and cooled (-10°C) mixture of 20.8 parts of carbon disulfide, 9 parts of N,N'-methanetetraylbis[cyclohexanamine] and 135 parts of tetrahydrofuran was added dropwise a solution of 15 parts of 4-[[1-(phenylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidineethanamine in tetrahydrofuran at a temperature below -10°C.
- 10 The reaction mixture was allowed to reach room temperature and the solvent was evaporated. The residue was crystallized from acetonitrile. The precipitate was filtered off and the filtrate was evaporated, yielding 15 parts (89%) of 2-[[1-(2-isothiocyanatoethyl)-4-piperidinyl]methyl]-1-(phenylmethyl)-1H-benzimidazole (395).
- 15 In a similar manner there were also prepared:



Comp. No.	R ¹	A ¹
396	4-F-C ₆ H ₄ -CH ₂	CH
397	2-thienylmethyl	CH
398	2-furanylmethyl	N
399	2-furanylmethyl	CH
400	4-F-C ₆ H ₄ -CH ₂	N
401	2-pyridinylmethyl	N
402	2-thienylmethyl	N
403	4-CH ₃ -C ₆ H ₄ -CH ₂	CH
404	4-CH ₃ -C ₆ H ₄ -CH ₂	N

Example 63

A mixture of 5.2 parts of 3,4-pyridinediamine, 19 parts of 2-[[1-(2-isothiocyanatoethyl)-4-piperidinyl)methyl]-1-[(4-methylphenyl)methyl]-1H-benzimidazole and 225 parts of tetrahydrofuran was stirred and refluxed overnight. The reaction mixture was evaporated, yielding 24 parts (100%) of N-(4-amino-3-pyridinyl)-N'-[2-[4-[[1-[(4-methylphenyl)methyl]-1H-benzimidazol-2-yl)methyl]-1-piperidinyl]-ethyl]thiourea (405).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:

- N-(4-amino-3-pyridinyl)-N'-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl)methyl]-1-piperidinyl]ethyl]thiourea (406);
- N-(4-amino-3-pyridinyl)-N'-[2-[4-[[1-(2-thienylmethyl)-1H-benzimidazol-2-yl)methyl]-1-piperidinyl]ethyl]thiourea (407);
- 15 N-(4-amino-3-pyridinyl)-N'-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1-piperidinyl]ethyl]thiourea (408);
- N-(4-amino-3-pyridinyl)-N'-[2-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl)methyl]-1-piperidinyl]ethyl]thiourea (409);
- N-(4-amino-3-pyridinyl)-N'-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1-piperidinyl]ethyl]thiourea (410);
- 20 N-(4-amino-3-pyridinyl)-N'-[2-[4-[[3-(2-pyridinylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1-piperidinyl]ethyl]thiourea (411);
- N-(4-amino-3-pyridinyl)-N'-[2-[4-[[3-(2-thienylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1-piperidinyl]ethyl]thiourea (412);
- 25 N-(4-amino-3-pyridinyl)-N'-[2-[4-[[1-(phenylmethyl)-1H-benzimidazol-2-yl)methyl]-1-piperidinyl]ethyl]thiourea (413); and
- N-(4-amino-3-pyridinyl)-N'-[2-[4-[[3-[(4-methylphenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1-piperidinyl]ethyl]thiourea (414).
- 30

Example 64

A mixture of 2 parts of N-(4-amino-3-pyridinyl)-N'-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl)methyl]-1-piperidinyl]-ethyl]thiourea, 3.3 parts of mercury(II) oxide, 0.1 parts of sulfur and 40 parts of ethanol was stirred and refluxed overnight. The

35

reaction mixture was filtered hot over Hyflo and the filtrate was evaporated. The residue was converted into the (E)-2-butenedioate salt in ethanol. The salt was filtered off and dried, yielding 1.8 parts (57%) of N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1H-imidazo[4,5-c]pyridin-2-amine (E)-2-butenedioate (1:3); mp. 184.7°C (415).

In a similar manner there were also prepared:

N-[2-[4-[[1-(2-thienylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1H-imidazo[4,5-c]pyridin-2-amine (E)-2-butenedioate (1:3). monohydrate; mp. 198.2°C (416);

N-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-1H-imidazo[4,5-c]pyridin-2-amine (E)-2-butenedioate (1:3). monohydrate; mp. 174.6°C (417);

N-[2-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1H-imidazo[4,5-c]pyridin-2-amine ethanedioate (1:4); mp. 201.1°C (418);

N-[2-[4-[[3-(2-pyridinylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-1H-imidazo[4,5-c]pyridin-2-amine ethanedioate (1:4); mp. 189.7°C (419);

N-[2-[4-[[3-(2-thienylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-1H-imidazo[4,5-c]pyridin-2-amine ethanedioate (2:5); mp. 154.5°C (420);

N-[2-[4-[[1-[(4-methylphenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1H-imidazo[4,5-c]pyridin-2-amine ethanedioate (1:3); mp. 203.5°C (421);

N-[2-[4-[[1-(phenylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1H-imidazo[4,5-c]pyridin-2-amine ethanedioate (1:4); mp. 199.0°C (422);

N-[2-[4-[[3-[(4-methylphenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-1H-imidazo[4,5-c]pyridin-2-amine ethanedioate (1:5); mp. 160.1°C (423); and

N-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-1H-imidazo[4,5-c]pyridin-2-amine ethanedioate (2:5); mp. 211.2°C (424).

Example 65

To a stirred and refluxing mixture of 3.7 parts of 4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1-piperidineethanamine and 90 parts of methylbenzene were added dropwise 1.1 parts of 2-pyridinecarboxyaldehyde using a water separator. Upon completion, stirring was continued for 20 hours at reflux. After cooling to 50°C, 44 parts of ethanol were added. At a temperature of 40°C, 0.4 parts of sodium borohydride were added portionwise. Upon completion, the whole was stirred for 2 hours at 45°C. The reaction mixture was poured into ice water and acetic acid while hot. The mixture was treated with ammonium hydroxide. The product was extracted three times with methylbenzene. The organic layer was washed with water, dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane, methanol and ammonium hydroxide (90:9:1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt in 2-propanone. The salt was filtered off and dried, yielding 3.0 parts (43.4%) of N-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1-piperidinyl]ethyl]-2-pyridinemethanamine (E)-2-butenedioate(1:2); mp. 160.7°C (425).

In a similar manner there were also prepared:

4-[[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl)methyl]-N-(phenylmethyl)-1-piperidineethanamine (426); and
N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl)methyl]-1-piperidinyl]ethyl]-2-pyridinemethanamine (E)-2-butenedioate(1:2); mp. 145.1°C (427).

Example 66

A mixture of 1.1 parts of isothiocyanatomethane, 5.5 parts of 4-[[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl)methyl]-1-piperidineethanamine and 90 parts of tetrahydrofuran was stirred overnight at room temperature. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions were collected and the eluent was

evaporated. The residue was converted into the ethanedioate salt in ethanol. The salt was filtered off and dried, yielding 4 parts (43%) of N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-N'-methylthiourea ethanedioate (1:2); mp.

5 169.0°C (428).

In a similar manner there were also prepared:

N-ethyl-N'-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyr-
idin-2-yl]-methyl]-1-piperidinyl]ethyl]urea; mp. 148.6°C (429); and

N-ethyl-N'-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]-
10 methyl]-1-piperidinyl]ethyl]urea; mp. 111.4°C (430).

Example 67

To a stirred mixture of 1.92 parts of 3-thiophenecarboxylic acid, 3.03 parts of N,N-diethylethanamine and 260 parts of dichloromethane were added 3.82 parts of 2-chloro-1-methylpyridinium iodide. The
15 whole was stirred for 1 hour and then 5.5 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidineethanamine were added. After stirring for 2 hours, water was added and the product was extracted with dichloromethane. The combined extracts
20 residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the
ethanedioate salt in ethanol and acetonitrile. The salt was filtered
25 off and crystallized from methanol. The product was filtered off and dried, yielding 3.5 parts (35.5%) of N-[2-[4-[[1-[(4-fluorophenyl)-methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-3-thiophene-
carboxamide ethanedioate(1:2); mp. 177.9°C (431).

In a similar manner there were also prepared:

30 N-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]-methyl]-1-piperidinyl]ethyl]-3-furancarboxamide; mp. 139.9°C (432);
N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1-methyl-1H-pyrrole-2-carboxamide ethanedioate
(2:5); mp. 164.9°C (433);

35

- N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-3-furancarboxamide ethanedioate(2:5).hemihydrate; mp. 139.7°C (434);
- 5 [2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]-methyl]-1-piperidinyl]ethyl]-3-pyridinecarboxylate ethanedioate (1:3); mp. 149.3°C (435);
- 3-amino-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]-methyl]-1-piperidinyl]ethyl]-2-pyrazinecarboxamide ethanedioate (1:2); mp. 166.8°C (436);
- 10 N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-4-thiazolecarboxamide ethanedioate(1:2); mp. 168.1°C (437);
- N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-methoxy-3-pyridinecarboxamide ethanedioate
- 15 (2:5); mp. 182.7°C (438);
- N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-9-oxo-2-thiopheneacetamide ethanedioate(1:2); mp. 180.2°C (439);
- N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-
- 20 piperidinyl]ethyl]-3-thiopheneacetamide ethanedioate(2:5); mp. 185.5°C (440);
- N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-methoxy-5-(1-oxobutyl)benzamide ethanedioate (1:2).hemihydrate; mp. 161.3°C (441);
- 25 N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1-methyl-1H-indole-2-carboxamide; mp. 137.3°C (442);
- N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-thiophenecarboxamide ethanedioate(1:2); mp.
- 30 157.6°C (443);
- N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-4-hydroxy-2-quinolinecarboxamide; mp. 262.4°C (444);
- N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-
- 35

piperidinyl]ethyl]-2-oxo-2H-1-benzopyran-3-carboxamide; mp. 134.0°C (445);

N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-thiazolecarboxamide ethanedioate(1:2); mp.

5 178.0°C (446); and

N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-4-pyridinecarboxamide ethanedioate(1:3) monohydrate; mp. 164.3°C (447).

Example 68

10 A mixture of 5.5 parts of 4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidineethanamine, 2.6 parts of 1,1'-carbonylbis[1H-imidazole] and 180 parts of tetrahydrofuran was stirred for 2 hours at room temperature. Gaseous N-methylmethan-

15 reaction : the cooler was provided with a CaCl₂ tube). The whole was stirred overnight at room temperature. The reaction mixture was evaporated. The residue was converted into the ethanedioate salt in 2-propanone. The precipitate was filtered off and the filtrate was evaporated. The residue was taken up in water. Ammonium hydroxide

20 was added till an alkaline solution was obtained. The product was extracted twice with methylbenzene. The combined extracts were washed with water, dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane, methanol and ammonium hydroxide (90:9:1 by volume)

25 as eluent. The pure and the less pure fractions were collected and the eluent was evaporated. The residue was purified by column chromatography (HPLC) over silica gel using a mixture of trichloromethane, methanol and methanol, saturated with ammonia, (96:3:1 by volume) as eluent. The pure fractions were collected and the eluent

30 was evaporated using a water bath at 30°C. The residue was converted into the ethanedioate salt in 2-propanone. The salt was allowed to crystallize while stirring. It was filtered off and dried in vacuo at 60°C, yielding 2.0 parts (20.0%) of N'-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]-

35 ethyl]-N,N-dimethylurea ethanedioate(2:5); mp. 94.2°C (448).

Example 69

A mixture of 25 parts of 1-[(4-fluorophenyl)methyl]-2-[[1-(2-isothiocyanatoethyl)-4-piperidinyl]methyl]-1H-benzimidazole and 160 parts of methanol saturated with ammonia was stirred overnight at room temperature. The reaction mixture was evaporated, yielding 25 parts (100%) of N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]thiourea (449).

Example 70

A mixture of 6.4 parts of methyl N-(2,2-dimethoxyethyl)-N'-methylcarbamimidothioate monohydroiodide, 7.3 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidine-ethanamine and 80 parts of 2-propanol was stirred and refluxed overnight. The reaction mixture was evaporated, yielding 12.77 parts (99%) of N-(2,2-dimethoxyethyl)-N'-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-N''-methylguanidine monohydroiodide (450).

In a similar manner there were also prepared:

N-(2,2-dimethoxyethyl)-N'-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]guanidine monohydroiodide (451).

Example 71

A mixture of 12.77 parts of N-(2,2-dimethoxyethyl)-N'-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-N''-methylguanidine monohydroiodide and 150 parts of a hydrochloric acid solution 10% was stirred and refluxed for 2 hours. Ice water was added and the whole was treated with a sodium hydroxide solution. The product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt in ethanol. The salt was filtered off and dried, yielding 3 parts (21%) of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-(1-methyl-1H-imidazol-2-yl)-1-piperidine-

ethanamine (E)-2-butenedioate (1:2) monohydrate; mp. 119.6°C (452).

In a similar manner there were also prepared:

4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-(1H-imidazol-2-yl)-1-piperidineethanamine ethanedioate(1:2).monohydrate
5 mp. 126.1°C (453).

Example 72

A mixture of 3.3 parts of 2-bromo-1-phenylethanone, 7 parts of N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]thiourea, 4 parts of potassium carbonate and 90
10 parts of tetrahydrofuran was stirred for 2 hours at room temperature. The reaction mixture was filtered over Hyflo and the filtrate was evaporated. The reaction mixture was poured into water. The product was extracted with dichloromethane. The extract was
15 washed with water, dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile. The
20 product was filtered off and dried, yielding 2.8 parts (33.3%) of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-(4-phenyl-2-thiazolyl)-1-piperidineethanamine; mp. 122.2°C (454).

In a similar manner there were also prepared:

ethyl 2-[[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]amino]-4-thiazolecarboxylate
25 ethanedioate(1:2); mp. 179.5°C (455); and 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-(4-methyl-2-thiazolyl)-1-piperidineethanamine ethanedioate(2:7); mp. 148.6°C (456).

30 Example 73

A mixture of 6 parts of ethyl 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidineacetate and 120 parts of a hydrochloric acid solution 6N was stirred and refluxed overnight. The reaction mixture was cooled and filtered. The filtrate was
35 evaporated and the semi-solid residue was taken up in about 120

parts of 2-propanol. The solution was filtered and about 70 parts of 2,2'-oxybispropane were added to the filtrate. After stirring for 2 hours at room temperature, the precipitated product was filtered off, dried overnight in vacuo at 80°C and pulverized in a mortar, yielding 3 parts (52%) of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl)methyl]-1-piperidineacetic acid dihydrochloride monohydrate; mp. 207.4°C (457).

Example 74

A mixture of 85 parts of 4-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl)methyl]-1-piperidinyl]-2-butanone and 600 parts of acetic acid was acidified with glacial acetic acid saturated with hydrogen bromide. A solution of 32.6 parts of bromine in acetic acid was added dropwise. Upon completion, stirring was continued overnight at room temperature. The reaction mixture was evaporated. The residue was stirred in 4-methyl-2-pentanone. The product was filtered off and dried, yielding 111 parts (80%) of 1-bromo-4-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl)methyl]-1-piperidinyl]-2-butanone trihydrobromide (458).

Example 75

A mixture of 0.75 parts of ethanethioamide, 7 parts of 1-bromo-4-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl)methyl]-1-piperidinyl]-2-butanone dihydrobromide and 80 parts of methanol was stirred overnight at room temperature. The reaction mixture was evaporated. Water was added. The whole was treated with sodium hydroxide. The product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (97:3 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was further purified by column chromatography (HPLC) over silica gel using a mixture of hexane, trichloromethane and methanol (45:45:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in ethanol and 2-propanone. The salt was filtered off and dried, yielding 2 parts

(33%) of 1-[(4-fluorophenyl)methyl]-2-[[1-[2-(2-methyl-4-thiazolyl)-ethyl]-4-piperidinyl]methyl]-1H-benzimidazole ethanedioate(2:5); mp. 124.1°C (459).

Example 76

- 5 To a stirred mixture of 5.5 parts of 4-[[1-[(4-fluorophenyl)-methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidineethanamine and 90 parts of tetrahydrofuran was added dropwise a solution of 3.8 parts of methyl 2-isothiocyanatobenzenecarboxylate in 18 parts of tetrahydrofuran (exothermic reaction). Upon completion, stirring was
- 10 continued for 1 hour. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from
- 15 acetonitrile. The product was filtered off and recrystallized from methanol. The product was filtered off and dried, yielding 3.6 parts (46%) of 3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]-methyl]-1-piperidinyl]ethyl]-2,3-dihydro-2-thioxo-4(1H)-quinazolinone; mp. 218.2°C (460).
- 20 In a similar manner there were also prepared:
- 3-[2-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2,3-dihydro-2-thioxo-4(1H)-quinazolinone; mp. 216.6°C (461);
- 3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2,3-dihydro-6-methyl-2-thioxothieno[2,3-d]-pyrimidin-4(1H)-one dihydrochloride. monohydrate; mp. 224.3°C (462);
- 25 3-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2,3-dihydro-2-thioxo-4(1H)-quinazolinone; mp. 204.2°C (463);
- 30 3-[2-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]-ethyl]-2,3-dihydro-6-methyl-2-thioxothieno[2,3-d]-pyrimidin-4(1H)-one; mp. 192.7°C (464); and
- 3-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2,3-dihydro-6-methyl-2-thioxothieno[2,3-d]-
- 35 pyrimidin-4(1H)-one; mp. 197.1°C (465).

Example 77

- To a stirred mixture of 4.9 parts of 2H-3,1-benzoxazine-2,4(1H)-dione and 45 parts of N,N-dimethylformamide were added dropwise 10.15 parts of 4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidineethanamine and 45 parts of C at 50°C. Upon completion, stirring was continued for 4 hours at 70°C. After cooling, the reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from 1,1'-oxybisethane. The product was filtered off and dried, yielding 10 parts (73%) of 2-amino-N-[2-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]benzamide; mp. 125.7°C (466).
- In a similar manner there were also prepared:
- N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-(methylamino)benzamide; mp. 84.3°C (467);
- 2-amino-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]benzamide; mp. 126.9°C (468);
- 2-amino-N-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]benzamide (469);
- N-[2-[4-[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2-(methylamino)benzamide ethane-dioate (2:5); mp. 172.3°C (470);
- 2-amino-N-[4-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]butyl]benzamide (471);
- 2-amino-N-[4-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]butyl]benzamide; mp. 127.7°C (472);
- 2-amino-N-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]benzamide; mp. 137.1°C (473);
- N-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2-(methylamino)benzamide; mp. 81.4°C (474); and
- 2-amino-N-[4-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]butyl]benzamide (475);
- N-[2-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piper-

idiny]l-ethyl]-2-(methylamino)benzamide (476);

2-amino-N-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-c]pyridin-2-yl]-methyl]-1-piperidinyl]-ethyl]benzamide ethanedioate(1:2); mp. 161.5°C (477);

- 5 2-amino-N-[2-[4-[[1-(2-thienylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]benzamide; mp. 143.5°C; (478)
- 2,3-dihydro-2,2-dimethyl-3-[2-[4-[[3-(2-pyridinylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-4(1H)-quinazolinone ethanedioate(1:1); mp. 210.2°C (479); and
- 10 2-amino-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]-oxy]-1-piperidinyl]ethyl]benzamide; mp. 141.7°C (480).

Example 78

- A mixture of 4 parts of 2-amino-N-[2-[4-[[1-[(4-fluorophenyl)-methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]benzamide,
- 15 20 parts of acetic acid anhydride and 40 parts of water was stirred and heated overnight at 120°C. After cooling, ice water was added. The whole was treated with ammonium hydroxide. The product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was converted into the (E)-2-butenedioate
- 20 salt in ethanol. The salt was filtered off and dried, yielding 3.7 parts (72%) of 3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-methyl-4(3H)-quinazolinone (E)-2-butenedioate (1:1); mp. 210.3°C (481).

- Following the same procedure and using equivalent amounts of
- 25 the appropriate starting materials, there were also prepared:
- 3-[2-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-methyl-4(3H)-quinazolinone trihydrochloride dihydrate; mp. 219.5°C (482);
- 3-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]-methyl]-1-piperidinyl]ethyl]-2-methyl-4(3H)-quinazolinone; mp.
- 30 147.6°C (483); and
- 3-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2-methyl-4(3H)-quinazolinone (E)-2-butenedioate (1:3); mp. 184.1°C (484).

Example 79

A mixture of 5 parts of 2-amino-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]oxy]-1-piperidinyl]ethyl]benzamide, 80 parts of 2-propanone and 1.9 parts of ethanedioic acid was stirred
5 for 1 hour at reflux temperature. After cooling, the product was filtered off and dried, yielding 4.8 parts (77%) of 3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]oxy]-1-piperidinyl]ethyl]-2,3-dihydro-2,2-dimethyl-4(1H)-quinazolinone ethanedioate (1:1); mp. 166.5°C (485).

10 Example 80

To a stirred mixture of 8 parts of N-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2-(methylamino)benzamide, 13 parts of N,N-diethylethanamine and 130 parts of dichloromethane was added dropwise a solution of 2.3 parts
15 of carbonylthioic dichloride in dichloromethane. Upon completion, stirring was continued for 2 hours. The reaction mixture was evaporated. The residue was crystallized from a mixture of methanol and ethanol. The product was filtered off and boiled in methanol. The product was filtered off and dried, yielding 3 parts (34.3%) of
20 3-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2,3-dihydro-1-methyl-2-thioxo-4(1H)-quinazolinone; mp. 169.2°C (486).

In a similar manner there were also prepared:

3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-
25 1-piperidinyl]ethyl]-2,3-dihydro-1-methyl-2-thioxo-4(1H)-quinazolinone; mp. 147.5°C (487); and
3-[2-[4-[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2,3-dihydro-1-methyl-2-thioxo-4(1H)-quinazolinone; mp. 176.1°C (488).

30 Example 81

A mixture of 10.3 parts of 2-amino-N-[4-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]-butyl]-benzamide, 3.2 parts of 1,1'-carbonylbis[1H-imidazole] and 180 parts of tetrahydrofuran was stirred and refluxed overnight. The
35 reaction mixture was evaporated. The residue was purified by column

chromatography over silica gel using first a mixture of trichloromethane and methanol saturated with ammonia (97:3 by volume) and then a mixture of trichloromethane and methanol saturated with ammonia (95:5 by volume) as eluent. The pure fractions were collected and the eluent was vaporated. The residue was crystallized from 2-propanone, yielding 3.8 parts (35.5%) of 3-[4-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]butyl]-2,4(1H,3H)-quinazolinedione; mp. 187.3°C (489).

Example 82

10 To a stirred mixture of 3.88 parts of 2-amino-N-[4-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]butyl]-benzamide, 2 parts of N,N-diethylethanamine and 90 parts of tetrahydrofuran were added dropwise 1.64 parts of trichloromethyl carbonochloridate. Upon completion, stirring was continued
15- overnight. Another portion of 1.6 parts of B was added and the whole was stirred overnight. The precipitate was filtered off and the filtrate was evaporated. Water was added to the residue. The solution was treated with ammonium hydroxide; The product was extracted with dichloromethane. The extract was dried, filtered and
20 evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 3 parts (73%) of 3-[4-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]butyl]-2,4(1H,3H)-quinazolinedione; mp. 185.5°C (490).

In a similar manner there was also prepared:

25 3-[4-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]butyl]-2,4-(1H,3H)-quinazolinedione; mp. 146.6°C (491).

Example 82

To a stirred mixture of 5.1 parts of 4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidineethanamine and 270
30 parts of tetrahydrofuran was added dropwise a solution of 3.8 parts of ethyl 2-isothiocyanatobenzoate in tetrahydrofuran. Upon completion, stirring was continued for 1 hour. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol,
35 saturated with ammonia, (97:3 by volume) as eluent. The pure

fractions were collected and the eluent was evaporated. The residue was crystallized from 2-propanone. The product was filtered off and dried, yielding 1.4 parts (18.6%) of 3-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2,3-dihydro-2-thioxo-4(1H)-quinazolinone; mp. 192.0°C (492).

Example 84

A mixture of 6 parts of N^1 -[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1,2-benzenediamine, 2.7 parts of 1,1'-thiocarbonylbis[1H-imidazole] and 90 parts of tetrahydrofuran was stirred and refluxed for 1 hour. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The oily residue was stirred in acetonitrile. The product was filtered off and crystallized from ethanol. The product was filtered off and dried, yielding 2.8 parts (41.5%) of 1-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1,3-dihydro-2H-benzimidazole-2-thione; mp. 157.1°C (493).

Example 85

A mixture of 7.5 parts of 6-chloro- N^4 -[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-4,5-pyrimidinediamine and 3.6 parts of urea was stirred and heated for 20 minutes at 220°C. Water was added to the reaction mixture. The precipitated product was filtered off and crystallized from methanol. The product was filtered off and recrystallized from a mixture of N,N -dimethylformamide and methanol, yielding 2.5 parts (32%) of 6-chloro-9-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-9H-purin-8-ol; mp. 243.0°C (494).

Example 86

A mixture of 11.3 parts of N^4 -[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-4,5-pyrimidinediamine, 3.75 parts of carbon disulfide and 117 parts of N,N -dimethylformamide was stirred overnight at room temperature. The

reaction mixture was poured into water. The product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 5 parts (40%) of 9-[2-[4-[[1-[(4-fluorophenyl)-methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-9H-purine-8-thiol; mp. 163.7°C (495).

Example 87

To a stirred mixture of 3 parts of 3-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-2,3-dihydro-6-methyl-2-thioxothieno[2,3-d]pyrimidin-4(1H)-one, 4.3 parts of potassium hydroxide, 56 parts of ethanol and 5.5 parts of water were added dropwise 45 parts of a hydrogen peroxide solution 3%. The whole was stirred overnight. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from 2-propanone. The product was filtered off and dried, yielding 1.7 parts (58%) of 3-[2-[4-[[3-(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]methyl]-1-piperidinyl]ethyl]-6-methylthieno[2,3-d]pyrimidine-2,4(1H,3H)-dione monohydrate; mp. 135.4°C (496).

In a similar manner there was also prepared:

3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-6-methyl-thieno[2,3-d]pyrimidine-2,4(1H,3H)-dione dihydrochloride. dihydrate; mp. 232.8°C (497).

Example 88

A mixture of 2.5 parts of 3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-methyl-4H-pyrido[1,2-a]pyrimidin-4-one and 120 parts of methanol was hydrogenated at normal pressure and at room temperature with 1 part of palladium-on-charcoal catalyst 10%. After the calculated amount

of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated. The residue was converted into the ethanedioate salt in ethanol. The salt was filtered off and dried, yielding 3 parts (87%) of 3-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-6,7,8,9-tetrahydro-2-methyl-4H-pyrido[1,2-a]pyrimidin-4-one ethanedioate (1:2); mp. 192.7°C (498).

Example 89

A mixture of 2.4 parts of 5-chloro-N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-pyridinamine, 1 part of calcium oxide and 120 parts of methanol was hydrogenated at normal pressure and at room temperature with 2 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated. Water was added to the residue. The product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was converted into the ethanedioate salt in a mixture of acetonitrile and ethanol. The salt was filtered off and dried in a dry pistol at 110-120°C, yielding 1.8 parts (50%) of N-[2-[4-(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-pyridinamine ethanedioate; mp. 156.2°C (499).

In a similar manner there were also prepared:

N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-3-pyridazinamine trihydrochloride monohydrate; mp. 197.9°C (500);

N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-4-pyrimidinamine; mp. 60.3°C (501);

9-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-9H-purin-8-ol; mp. 213.6°C (502);

N⁴-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-4,5-pyrimidinediamine (503).

Example 90

A mixture of 7.7 parts of 1-[(4-fluorophenyl)methyl]-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1H-benzimidazole and

150 parts of a hydrobromic acid solution 48% in water was stirred overnight at 80°C. The reaction mixture was evaporated and water was added to the residue. The whole was treated with ammonium hydroxide and the product was extracted with trichloromethane. The
5 extract was dried, filtered and evaporated. The residue was purified by column chromatography (HPLC) over silica gel using a mixture of hexane, trichloromethane and methanol (45:45:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile. The
10 product was filtered off and dried, yielding 2.5 parts (35%) of 4-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]phenol; mp. 130.3°C (504).

In a similar manner there were also prepared:

1-[(4-fluorophenyl)methyl]-2-[[1-[2-(4-hydroxyphenyl)ethyl]-4-piperidinyl]-methyl]-1H-benzimidazol-6-ol monohydrate; mp. 169.4°C
15 (505).

Example 91

A mixture of 7 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-(2-nitrophenyl)-1-piperidineethanamine,
20 1 part of a solution of thiophene in methanol 4% and 200 parts of methanol was hydrogenated at normal pressure and at 50°C with 2 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated, yielding 6 parts (90%) of N¹-[2-[4-
25 [[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-1,2-benzenediamine (506).

Example 92

To a stirred mixture of 34.5 parts of 9-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]-ethyl]-9H-purine-8-thiol and 180 parts of N,N-dimethylformamide were
30 added portionwise 3.2 parts of a sodium hydride dispersion 50%. Upon completion, stirring was continued for 0.5 hours at room temperature. 11.5 Parts of iodomethane were added dropwise. After complete addition, the whole was stirred for 1 hour. The reaction
35 mixture was poured into water. The product was extracted with

4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 28.3 parts (80%) of 9-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-8-(methylthio)-9H-purine; mp. 133.1°C (507).

Example 93

To a stirred and cooled mixture of 6.2 parts of N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-pyridinamine, 2 parts of N,N-diethylethanamine and 90 parts of tetrahydrofuran was added dropwise slowly a solution of 1.9 parts of benzoyl chloride in 45 parts of tetrahydrofuran. Upon completion, stirring was continued for 2 hours. The reaction mixture was evaporated. Water was added to the residue. The solution was treated with ammonium hydroxide. The product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was crystallized from 2,2'-oxybispropane. The product was filtered off and dried, yielding 5.3 parts (69%) of N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-N-(2-pyrimidinyl)benzamide; mp. 108.1°C (508).

In a similar manner there were also prepared: N-[2-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-N-(2-pyrimidinyl)-2-furancarboxamide ethanedioate (1:2); mp. 147.7°C (509).

Example 94

A mixture of 6.6 parts of N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-2-pyrimidinamine, 20 parts of acetic acid anhydride and 60 parts of water was stirred and refluxed overnight. The reaction mixture was evaporated. Water was added and the whole was treated with ammonium hydroxide. The product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in 2-propanone. The

salt was filtered off and dried, yielding 2.7 parts (27%) of
N-[2-[4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-
piperidinyl]ethyl]-N-(2-pyrimidinyl)acetamide ethanedioate (1:2);
mp. 173.7°C (510).

5 Example 95

A mixture of 2.9 parts of N-[2-[4-[[1-(2-furanylmethyl)-1H-
benzimidazol-2-yl]methyl]-1-piperidinyl]-ethyl]-2-(methyl-
amino)benzamide, 10 parts of acetic acid anhydride and 20 parts of
10 water was stirred and heated for 3 hours at 100°C. The reaction
mixture was cooled, water was added and the whole was made alkaline
with ammonium hydroxide. The product was extracted with dichloro-
methane. The extract was dried, filtered and evaporated. The residue
was purified twice by column chromatography over silica gel using
each time a mixture of trichloromethane and methanol, saturated with
15 ammonia (95:5 by volume) as eluent. The pure fractions were
collected and the eluent was evaporated. The residue was converted
into the ethanedioate salt in ethanol, yielding 0.2 parts (4.3%) of
2-(acetylmethylamino)-N-[2-[4-[[1-(2-furanylmethyl)-1H-benzimidazol-
2-yl]-methyl]-1-piperidinyl]ethyl]benzamide ethanedioate (2:5);
20 mp. 146.8°C (511).

Example 96

A mixture of 13.4 parts of 4-[[1-[(4-fluorophenyl)methyl]-1H-benz-
imidazol-2-yl]methyl]-N-(phenylmethyl)-1-piperidineethanamine, 4
parts of poly(oxymethylene), 1 part of a solution of thiophene in
25 methanol 4% and 120 parts of methanol was hydrogenated at normal
pressure and at room temperature with 2 parts of palladium-on-
charcoal catalyst 5%. After the calculated amount of hydrogen was
taken up, the catalyst was filtered off and the filtrate was
evaporated. The residue was taken up in 4-methyl-2-pentanone. The
30 solution was washed with water, dried, filtered and evaporated. The
residue was converted into the ethanedioate salt in methanol. The
salt was filtered off and dried, yielding 13.02 parts (68.3%) of
4-[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-methyl-
N-(phenylmethyl)-1-piperidine-ethanamine ethanedioate (1:2);
35 mp. 172.6°C (512).

A mixture of 10 parts of 4-[[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-methyl-N-(phenylmethyl)-1-piperidine-ethanamine and 120 parts of methanol was hydrogenated at normal pressure and at room temperature with 2 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated. The residue was converted into the ethanedioate salt in methanol. The salt was filtered off and dried, yielding 7 parts (87.8%) of 4-[[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-methyl-1-piperidineethanamine ethanedioate(1:2); mp. 205.8°C (dec.) (513).

A mixture of 1.8 parts of 2-chloropyrimidine, 6 parts of 4-[[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-N-methyl-1-piperidineethanamine, 1.7 parts of sodium hydrogen carbonate and 120 parts of ethanol was stirred and refluxed overnight. The reaction mixture was evaporated. Water was added. The product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was crystallized from a mixture of 2,2'-oxybispropane and 1,1'-oxybisethane (50:50 by volume). The product was filtered off and dried, yielding 5.5 parts (76.5%) of N-[2-[4-[[[1-[(4-fluorophenyl)methyl]-1H-benzimidazol-2-yl]methyl]-1-piperidinyl]ethyl]-N-methyl-2-pyrimidinamine; mp. 135.4°C (514).

Example 97

To a stirred mixture of 3.5 parts of 2-[[[1-[2-(4-methoxyphenyl)-ethyl]-4-piperidinyl]methyl]-1H-benzimidazole and 18 parts of N,N-dimethylacetamide were added portionwise 0.5 parts of a sodium hydride dispersion 59.4% at room temperature. After stirring for 35 minutes at room temperature and for 10 minutes at 60°C, a solution of 1.7 parts of 1-(chloromethyl)-2-fluorobenzene in 9 parts of N,N-dimethylacetamide was added dropwise at 60°C. After stirring for 10 minutes, the reaction mixture was cooled and poured into 150 parts of water. The product was extracted twice with 4-methyl-2-pentanone. The combined extracts were washed with water, dried, filtered and evaporated. The residue was crystallized from a mixture of 2,2'-oxybispropane and 2-propanone. The product was filtered off

and dried, yielding 3.0 parts (65.5%) of 1-[(2-fluorophenyl)methyl]-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1H-benzimidazole; mp. 109.3°C (515).

In a similar manner there were also prepared:


- 5 2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1-(2-phenylethyl)-1H-benzimidazole dihydrochloride monohydrate; mp. 176.0°C (516);
- 1-(diphenylmethyl)-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1H-benzimidazole ethanedioate(2:5); mp. 174.0°C (517);
- 10 1-[(2,5-dimethylphenyl)methyl]-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1H-benzimidazole; mp. 118.3°C (518);
- 1-[(2,6-dichlorophenyl)methyl]-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1H-benzimidazole; mp. 152.4°C (519);
- 1-[(3-chlorophenyl)methyl]-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1H-benzimidazole dihydrochloride; mp. 173.3°C (520);
- 15 2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1-(1-naphthalenylmethyl)-1H-benzimidazole ethanedioate(2:5); mp. 186.8°C (521);
- 1-cyclohexyl-2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1H-benzimidazole dihydrochloride monohydrate; mp. 189.6°C (522);
- 20 2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1-(3-thienylmethyl)-1H-benzimidazole ethanedioate(1:2); mp. 185.5°C (523);
- 2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1-(2-pyrazinylmethyl)-1H-benzimidazole (E)-2-butenedioate(1:1); mp. 180.9°C (524);
- 2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1-[(5-methyl-2-thienyl)methyl]-1H-benzimidazole ethanedioate(1:2); mp. 194.9°C (525); and
- 25 2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]methyl]-1-[(3-methyl-2-thienyl)methyl]-1H-benzimidazole ethanedioate(1:2) monohydrate; mp. 166.2°C (526).

30

35

Example 98


Following the procedures described in example 18 there were also prepared:



N°	R ^{1-a}	A ¹ =A ² -A ³ =A ⁴
527	4-F-C ₆ H ₄ -	CH=CH-CH=C(OCH ₃)
528	4-F-C ₆ H ₄ -	C(OCH ₃)=CH-CH=CH
529	4-F-C ₆ H ₄ -	CH=C(OCH ₃)-C(OCH ₃)=CH
530	4-F-C ₆ H ₄ -	CH=C(OCH ₃)-CCl=CH
531	4-F-C ₆ H ₄ -	CH=CCl-C(OCH ₃)=CH
532	2-furanyl	CH=CH-C(OCH ₃)=CH
533	2-furanyl	CH=C(OCH ₃)-CH=CH
534	2-pyridinyl	CH=CH-C(OCH ₃)=CH
535	2-pyridinyl	CH=C(OCH ₃)-CH=CH
536	4-F-C ₆ H ₄ -	N=C(OCH ₃)-CH=CH
537	2-furanyl	N=C(OCH ₃)-CH=CH
538	2-pyridinyl	N=C(OCH ₃)-CH=CH

Example 99

Following the procedures described in example 26 there were also prepared:



10


N°	R ^{1-a}	A ¹ =A ² -A ³ =A ⁴
539	4-F-C ₆ H ₄ -	CH=CH-CH=C(OCH ₃)
540	4-F-C ₆ H ₄ -	C(OCH ₃)=CH-CH=CH
541	4-F-C ₆ H ₄ -	CH=C(OCH ₃)-C(OCH ₃)=CH
542	4-F-C ₆ H ₄ -	CH=C(OCH ₃)-CCl=CH
543	4-F-C ₆ H ₄ -	CH=CCl-C(OCH ₃)=CH
544	2-furanyl	CH=CH-C(OCH ₃)=CH
545	2-furanyl	CH=C(OCH ₃)-CH=CH
546	2-pyridinyl	CH=CH-C(OCH ₃)=CH
547	2-pyridinyl	CH=C(OCH ₃)-CH=CH
548	4-F-C ₆ H ₄ -	N=C(OCH ₃)-CH=CH
549	2-furanyl	N=C(OCH ₃)-CH=CH
550	2-pyridinyl	N=C(OCH ₃)-CH=CH

15

20

Example 100

25 Following the procedures described in example 34 there were also prepared:




35

N°	R ^{1-a}	A ¹ =A ² -A ³ =A ⁴
551	4-F-C ₆ H ₄ -	CH=CH-CH=C(OH)
552	4-F-C ₆ H ₄ -	C(OH)=CH-CH=CH
553	4-F-C ₆ H ₄ -	CH=C(OH)-C(OH)=CH

554	4-F-C ₆ H ₄ -	CH=C(OH)-CCl=CH
555	4-F-C ₆ H ₄ -	CH=CCl-C(OH)=CH
556	2-furanyl	CH=CH-C(OH)=CH
557	2-furanyl	CH=C(OH)-CH=CH
558	2-pyridinyl	CH=CH-C(OH)=CH
559	2-pyridinyl	CH=C(OH)-CH=CH
560	4-F-C ₆ H ₄ -	N=C(OH)-CH=CH
561	2-furanyl	N=C(OH)-CH=CH
562	2-pyridinyl	N=C(OH)-CH=CH

Example 101

Following the procedures described in example 46 there were also prepared:



N°	Ar	R ^{1-a}	A ¹ =A ² -A ³ =A ⁴
563	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=CH-CH=C(OH)
564	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=CH-C(OH)=CH
565	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	C(OH)=CH-CH=CH
566	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=C(OH)-C(OH)=CH
567	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=CCl-C(OH)=CH
568	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=C(OH)-CCl=CH
569	3,4-(CH ₃ O) ₂ -C ₆ H ₃ -	4-F-C ₆ H ₄ -	CH=CH-C(OH)=CH
570	3,4-(CH ₃ O) ₂ -C ₆ H ₃ -	4-F-C ₆ H ₄ -	CH=C(OH)-CH=CH
571	3-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=CH-C(OH)=CH
572	3-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=C(OH)-CH=CH
573	4-CH ₃ O-C ₆ H ₄ -	2-furanyl	CH=CH-C(OH)=CH
574	4-CH ₃ O-C ₆ H ₄ -	2-furanyl	CH=C(OH)-CH=CH

5

10

15

20


25

30

575	4-CH ₃ O-C ₆ H ₄ -	2-pyridinyl	CH=CH-C(OH)=CH
576	4-CH ₃ O-C ₆ H ₄ -	2-pyridinyl	CH=C(OH)-CH=CH
577	4-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	N=CH-CH=CH
578	4-CH ₃ O-C ₆ H ₄ -	2-pyridinyl	N=CH-CH=CH
579	4-CH ₃ O-C ₆ H ₄ -	2-furanyl	N=CH-CH=CH
580	4-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	N=C(OH)-CH=CH
581	4-CH ₃ O-C ₆ H ₄ -	2-pyridinyl	N=C(OH)-CH=CH
582	4-CH ₃ O-C ₆ H ₄ -	2-furanyl	N=C(OH)-CH=CH
583	2-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=CH-C(OH)=CH
584	2-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=C(OH)-CH=CH
585	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=CH-CH=C(OCH ₃)
586	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=CH-C(OCH ₃)=CH
587	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	C(OCH ₃)=CH-CH=CH
588	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=C(OCH ₃)-C(OCH ₃)=CH
589	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=CCl-C(OCH ₃)=CH
590	4-CH ₃ O-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=C(OCH ₃)-CCl=CH
591	3,4-(CH ₃ O) ₂ -C ₆ H ₃ -	4-F-C ₆ H ₄ -	CH=CH-C(OCH ₃)=CH
592	3,4-(CH ₃ O) ₂ -C ₆ H ₃ -	4-F-C ₆ H ₄ -	CH=C(OCH ₃)-CH=CH
593	3-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=CH-C(OCH ₃)=CH
594	3-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=C(OCH ₃)-CH=CH
595	4-CH ₃ O-C ₆ H ₄ -	2-furanyl	CH=CH-C(OCH ₃)=CH
596	4-CH ₃ O-C ₆ H ₄ -	2-furanyl	CH=C(OCH ₃)-CH=CH
597	4-CH ₃ O-C ₆ H ₄ -	2-pyridinyl	CH=CH-C(OCH ₃)=CH
598	4-CH ₃ O-C ₆ H ₄ -	2-pyridinyl	CH=C(OCH ₃)-CH=CH
599	4-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	N=C(OCH ₃)-CH=CH
600	4-CH ₃ O-C ₆ H ₄ -	2-pyridinyl	N=C(OCH ₃)-CH=CH
601	4-CH ₃ O-C ₆ H ₄ -	2-furanyl	N=C(OCH ₃)-CH=CH
602	2-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=CH-C(OCH ₃)=CH
603	2-CH ₃ O-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=C(OCH ₃)-CH=CH

In a similar manner there were also prepared:

35



5

10








15

20

25

30

35

N°	L'	n	A ¹ =A ² -A ³ =A ⁴
604		2	CH=C(OH)-CH=CH
605		2	CH=CH-C(OH)=CH
606		2	CH=C(OH)-C(OH)=CH
607		3	CH=C(OH)-CH=CH
608		3	CH=CH-C(OH)=CH
609		3	CH=C(OH)-C(OH)=CH
610		3	CH=C(OH)-CH=CH






5

10

15

20

25

611		3	$\text{CH}=\text{CH}-\text{C}(\text{OH})=\text{CH}$
612		3	$\text{CH}=\text{C}(\text{OH})-\text{C}(\text{OH})=\text{CH}$
613	2-thienyl	2	$\text{CH}=\text{C}(\text{OH})-\text{CH}=\text{CH}$
614	2-thienyl	2	$\text{CH}=\text{CH}-\text{C}(\text{OH})=\text{CH}$
615	2-thienyl	2	$\text{CH}=\text{C}(\text{OH})-\text{C}(\text{OH})=\text{CH}$
616		2	$\text{CH}=\text{C}(\text{OH})-\text{CH}=\text{CH}$
617		2	$\text{CH}=\text{CH}-\text{C}(\text{OH})=\text{CH}$
618		2	$\text{CH}=\text{C}(\text{OH})-\text{C}(\text{OH})=\text{CH}$


30

Example 102

Following the procedures described in example 90 there were also prepared:

35

5



10

15

20

25

30

N°	Ar	R ^{1-a}	A ¹ =A ² -A ³ =A ⁴
619	4-HO-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=CH-CH=C(OH)
620	4-HO-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=CH-C(OH)=CH
621	4-HO-C ₆ H ₄	4-F-C ₆ H ₄ -	C(OH)=CH-CH=CH
622	4-HO-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=C(OH)-C(OH)=CH
623	4-HO-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=CCl-C(OH)=CH
624	4-HO-C ₆ H ₄	4-F-C ₆ H ₄ -	CH=C(OH)-CCl=CH
625	3,4-(HO) ₂ -C ₆ H ₃ -	4-F-C ₆ H ₄ -	CH=CH-C(OH)=CH
626	3,4-(HO) ₂ -C ₆ H ₃ -	4-F-C ₆ H ₄ -	CH=C(OH)-CH=CH
627	3-HO-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=CH-C(OH)=CH
628	3-HO-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=C(OH)-CH=CH
629	4-HO-C ₆ H ₄ -	2-furanyl	CH=CH-C(OH)=CH
630	4-HO-C ₆ H ₄ -	2-furanyl	CH=C(OH)-CH=CH
631	4-HO-C ₆ H ₄ -	2-pyridinyl	CH=CH-C(OH)=CH
632	4-HO-C ₆ H ₄ -	2-pyridinyl	CH=C(OH)-CH=CH
633	4-HO-C ₆ H ₄ -	4-F-C ₆ H ₄ -	N=CH-CH=CH
634	4-HO-C ₆ H ₄ -	2-pyridinyl	N=CH-CH=CH
635	4-HO-C ₆ H ₄ -	2-furanyl	N=CH-CH=CH
636	4-HO-C ₆ H ₄ -	4-F-C ₆ H ₄ -	N=C(OH)-CH=CH
637	4-HO-C ₆ H ₄ -	2-pyridinyl	N=C(OH)-CH=CH
638	4-HO-C ₆ H ₄ -	2-furanyl	N=C(OH)-CH=CH
639	2-HO-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=CH-C(OH)=CH
640	2-HO-C ₆ H ₄ -	4-F-C ₆ H ₄ -	CH=C(OH)-CH=CH

35

The useful antihistaminic properties of the compounds of formula (I) are demonstrated in the following test procedure.

Protection of rats from compound 48/80-induced lethality.

5

Compound 48/80, a mixture of oligomers obtained by condensation of 4-methoxy-N-methylbenzeneethanamine and formaldehyde has been described as a potent histamine releasing agent (Int. Arch. Allergy, 13, 336 (1958)). The protection from compound 48/80-induced lethal circulatory collapse appears to be a simple way of evaluating quantitatively the antihistaminic activity of test compounds. Male rats of an inbred Wistar strain, weighing 240-260 g were used in the experiment. After overnight starvation the rats were transferred to conditioned laboratories (temp. = $21 \pm 1^\circ\text{C}$, relative humidity = $65 \pm 5\%$).

15

The rats were treated subcutaneously or orally with a test compound or with the solvent (NaCl solution, 0.9%). One hour after treatment there was injected intravenously compound 48/80, freshly dissolved in water, at a dose of 0.5 mg/kg (0.2 ml/100 g of body weight). In control experiments, wherein 250 solvent-treated animals were injected with the standard dose of compound 48/80, not more than 2.8% of the animals survived after 4 hours. Survival after 4 hours is therefore considered to be a safe criterion of a protective effect of drug administration.

20

The ED_{50} -values of the compounds of formula (I) are listed in the first column of tables 1 to 4. Said ED_{50} -values are the values in mg/kg body weight at which the tested compounds protect 50% of the tested animals against compound 48/80-induced lethality.

25

The compounds of formula (I) and the pharmaceutically acceptable acid addition salts thereof are also potent serotonin-antagonists. The potency of the subject compounds as serotonin-antagonists is clearly evidenced by the results obtained in the following tests wherein the antagonistic activity of the subject compounds on the effect of serotonin is examined.

30

35

Antagonistic activity in the effects of serotonin in the gastric lesion test.

5 A. Lesions induced by compound 48/80:

Compound 48/80 (a mixture of oligomers obtained by condensation of 4-methoxy-N-methylbenzeneethanamine and formaldehyde) is a potent releaser of vasoactive amines from endogenous stores such as, for example, histamine and serotonin. Rats injected with compound 10 48/80 exhibit consistent changes of blood flow in different vascular beds: cyanosis of the ears and the extremities are prominent within five minutes after injection of the compound; the rats die from shock within 30 minutes. The shock, followed by death, can be avoided if the rats are pretreated with a classical H₁ 15 antagonist.

However the stimulatory effects on gastric secretion are not suppressed so that rats treated with compound 48/80 and protected from shock by an H₁ antagonist may exhibit all signs of intensive gastric gland activity: gross autopsy shows distended stomachs with 20 abnormal contents and rough bright red patches all over the mucosa, corresponding to areas of disintegrated glands. A number of known serotonin-antagonists such as, for example, methysergide, cyproheptadine; cinanserin, mianserin, pipamperone, spiperone, pizotifen and metergoline, prevent completely the cyanosis of ears and 25 extremities as well as the lesions in the glandular area of the stomach and the abnormal gastric distension.

B. Method:

Male rats of a Wistar inbred strain, weighing 220-250 g, were starved overnight, water being available ad libitum. The test 30 compounds were administered orally as a solution or as a suspension in aqueous medium. A control rat and a "blank" rat received the test compound. One hour later 5-[4-(diphenylmethyl)-1-piperazinylmethyl]-1-methyl-1H-benzimidazole-2-methanol was administered subcutaneously to all rats at the dose of 2.5 mg/kg. Two hours after 35 the oral or subcutaneous administration of the test compound,

the compound 48/80 (freshly solved in water at a concentration of 0.25 mg/ml) was injected intravenously into all rats (dose: 1 mg/kg) except the "blank" rats.

- Four hours after the intravenous injection of compound 48/80, the rats were decapitated and the stomachs were removed. Subsequently the stomachs were inspected for distension and contents (blood, fluid, food) and thoroughly rinsed. The macroscopic lesions were scored from 0 to +++, 0 corresponding to complete absence of visible lesions and the highest score corresponding to reddish rough patches covering more than half the glandular area.

The second column of tables 1 - 4 show for a number of compounds of formula (I) the doses (in mg/kg body weight) at which the distension of the stomach as well as the lesions in the glandular area of the stomach are completely absent in 50% of the test rats

- (ED₅₀-value).

The compounds listed in table 1 are not given for the purpose of limiting the invention thereto but only to exemplify the useful pharmacological activities of all the compounds within the scope of formula (I).

20

25

30

35

Compound 48/80 gastric lesion .		
Comp. lethality test in test		
No.	rats ED ₅₀ in	ED ₅₀ in mg/kg
	mg/kg body weight	body weight
8	0.04	-
23	0.08	-
25	0.16	-
34	0.04	0.63
40	0.08	-
42	0.02	-
45	0.04	-
47	0.08	0.63
48	0.16	-
50	0.04	-
59	0.02	-
64	0.04	-
66	0.02	-
81	0.08	-
82	0.005	0.63
83	0.01	-
85	0.01	-
86	0.01	-
89	0.04	-
90	0.01	0.31
93	0.04	-
94	0.08	0.63
96	0.04	-
98	0.04	-
99	0.04	0.31
100	0.16	-
101	0.02	0.31
102	0.02	0.31

Compound 48/80 gastric lesion		
Comp. lethality test in test		
N .	rats ED ₅₀ in	ED ₅₀ in mg/kg
	mg/kg body weight	body weight
103	0.005	0.16
104	0.08	-
105	0.01	-
107	0.04	0.08
110	0.02	-
112	0.08	-
113	0.08	-
115	0.16	-
116	0.16	-
117	0.01	-
118	0.04	0.63
123	0.01	-
124	0.04	0.31
126	0.04	0.63
127	0.08	0.63
128	0.16	0.63
129	0.16	-
130	0.16	-
131	0.02	0.16
133	0.02	0.16
134	0.04	-
139	0.08	-
140	0.04	-
142	0.02	-
143	0.04	0.63
144	0.04	-

145	0.02	0.16
147	0.08	-
149	0.04	-
151	0.02	0.16
152	0.04	-
154	0.08	-
158	0.04	-
161	0.04	-
162	0.08	0.31
163	0.02	-
164	0.02	0.02
165	0.02	-
166	0.04	-
167	0.01	-
168	0.04	0.63
169	0.02	-
170	0.08	-
171	0.08	-
173	0.01	-
174	0.01	-
175	0.02	-
176	0.04	-
177	0.04	0.31
178	0.02	-
180	0.01	0.31
181	0.04	-
182	0.04	0.63
183	0.01	-
184	0.02	0.63
186	0.04	0.31
187	0.04	0.31
189	0.02	0.63
198	0.04	0.08

202	0.08	-
203	0.04	-
204	0.08	-
205	0.08	-
207	0.08	0.08
208	0.04	0.08
209	0.08	0.63
214	0.08	0.04
215	0.08	-
216	0.04	-
218	0.04	0.04
219	0.08	0.63
221	0.04	-
224	0.04	0.08
226	0.08	0.04
228	0.04	0.01
229	0.04	0.63
232	0.08	-
234	0.02	-
236	0.08	0.63
238	0.08	0.63
243	0.04	-
244	0.16	0.31
245	0.02	-
248	0.04	-
255	0.16	-
256	0.08	-
257	0.02	-
258	0.08	-
259	0.08	-
260	0.01	0.63
261	0.08	-
262	0.04	0.16

0151826

263	0.04	0.04
264	0.04	0.63
265	0.02	-
266	0.04	-
268	0.16	-
270	0.08	-
272	0.08	0.63
273	0.01	0.31
274	0.16	-
276	0.04	0.31
283	0.04	0.63
290	0.08	-
293	0.08	-
297	0.08	-
328	0.16	-
329	0.31	-
330	0.08	0.16
331	0.04	0.02
332	0.08	0.08
333	0.04	-
334	0.005	0.31
335	0.02	0.16
336	0.01	0.08
338	0.02	-
339	0.01	-
340	0.08	-
342	0.02	0.16
344	0.02	-
345	0.08	-
347	0.08	0.63
348	0.04	0.04
349	0.08	-
351	0.02	0.08

352	0.01	-
355	0.08	-
356	0.02	0.08
357	0.02	0.63
361	0.02	-
362	0.08	0.63
363	0.04	-
364	0.04	-
365	0.08	-
366	0.31	-
367	0.16	0.16
368	0.01	-
369	0.04	-
370	0.02	-
371	0.16	0.63
372	0.04	0.16
373	0.08	-
374	0.02	0.02
376	0.02	0.63
377	0.16	-
378	0.04	-
379	0.08	0.31
380	0.31	0.63
381	0.08	0.16
382	0.08	0.63
383	0.04	0.16
384	0.08	0.63
385	0.01	0.08
388	0.16	-
390	0.04	-
391	0.02	0.02
392	0.02	0.63
393	0.08	-

394	0.02	-
397	0.04	0.04
399	0.02	0.04
402	0.04	-
416	0.04	0.04
417	0.04	-
418	0.02	0.04
422	0.04	0.04
427	0.16	-
428	0.04	0.01
429	0.02	-
430	0.01	0.005
431	0.04	0.005
432	0.04	0.31
433	0.01	0.16
434	0.0025	-
436	0.01	0.63
438	0.08	0.16
439	0.16	0.31
440	0.02	-
441	0.08	0.16
443	0.02	0.16
444	0.16	-
445	0.16	-
448	0.08	0.16
452	0.16	-
453	0.04	0.02
455	0.04	0.63
456	0.01	0.63
459	0.02	-
460	0.08	0.63
461	0.08	0.08
463	0.08	-

466	0.02	-
467	0.04	0.08
468	0.02	0.63
470	0.04	-
472	0.04	-
473	0.02	-
477	0.02	0.63
478	0.02	0.04
479	0.04	-
480	0.02	-
489	0.08	0.63
490	0.31	-
491	0.16	-
493	0.08	0.04
495	0.04	0.16
498	0.08	0.16
499	0.04	0.01
500	0.01	-
501	0.04	-
502	0.08	0.16
504	0.08	0.63
505	0.04	0.04
507	0.16	-
508	0.08	0.16
509	0.02	-
510	0.02	0.16
511	0.16	-
512	0.08	-
513	0.08	-
514	0.02	0.63
515	0.16	-
516	0.16	-
523	0.04	0.31
524	0.16	-

In view of their antihistaminic and serotonin-antagonistic properties, the compounds of formula (I) and their acid-addition salts are very useful in the treatment of allergic diseases such as, for example, allergic rhinitis, allergic conjunctivities, chronic
5 urticaria, allergic asthma and the like.

In view of their useful antihistaminic and serotonin-antagonistic activity, the subject compounds may be formulated into various pharmaceutical forms for administration purposes.
To prepare the pharmaceutical compositions of this invention, an
10 effective amount of the particular compound, in base or acid-addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration.

15 These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils,
20 alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules
25 represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may
30 be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed.

In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wettable agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment.

Acid addition salts of (I) due to their increased water solubility over the corresponding base form, are obviously more suitable in the preparation of aqueous compositions.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

"Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I), a possible stereochemically isomeric form or pharmaceutically acceptable acid addition salt thereof.

Example 98 : ORAL DROPS

500 Grams of the A.I. was dissolved in 0.5 liters of 2-hydroxypropanoic acid and 1.5 liters of the polyethylene glycol at 60-80°C. After cooling to 30-40°C there were added 35 liters of polyethylene glycol and the mixture was stirred well. Then there was added a solution of 1750 grams of sodium saccharin in 2.5 liters of purified water and while stirring there were added 2.5 liters of cocoa flavor and polyethylene glycol q.s. to a volume of 50 liters, providing an

oral drop solution comprising 10 milligrams of the A.I. per milliliter. The resulting solution was filled into suitable containers.

Example 99 : ORAL SOLUTION

9 Grams of methyl 4-hydroxybenzoate and 1 gram of propyl
5 4-hydroxybenzoate were dissolved in 4 liters of boiling purified water. In 3 liters of this solution were dissolved first 10 grams of 2,3-dihydroxybutanedioic acid and thereafter 20 grams of the A.I. The latter solution was combined with the remaining part of the former solution and 12 liters 1,2,3-propanetriol and 3 liters of sorbitol 70%
10 solution were added thereto. 40 Grams of sodium saccharin were dissolved in 0.5 liters of water and 2 milliliters of raspberry and 2 milliliters of gooseberry essence were added. The latter solution was combined with the former, water was added q.s. to a volume of 20 liters providing an oral solution comprising 20 milligrams of the
15 active ingredient per teaspoonful (5 milliliters). The resulting solution was filled in suitable containers.

Example 100 : CAPSULES

20 Grams of the A.I., 6 grams sodium lauryl sulfate, 56 grams starch, 56 grams lactose, 0.8 grams colloidal silicon dioxide, and 1.2
20 grams magnesium stearate were vigorously stirred together. The resulting mixture was subsequently filled into 1000 suitable hardened gelating capsules, comprising each 20 milligrams of the active ingredient.

Example 101 : FILM-COATED TABLETS

25 Preparation of tablet core

A mixture of 100 grams of the A.I., 570 grams lactose and 200 grams starch was mixed well and thereafter humidified with a solution of 5 grams sodium dodecyl sulfate and 10 grams polyvinylpyrrolidone in about 200 milliliters of water. The wet powder mixture was sieved,
30 dried and sieved again. Then there was added 100 grams microcrystalline cellulose and 15 grams hydrogenated vegetable oil. The whole was mixed well and compressed into tablets, giving 10,000 tablets, each containing 10 milligrams of the active ingredient.

Coating

35 To a solution of 10 grams methyl cellulose in 75 milliliters of

denaturated ethanol there was added a solution of 5 grams of ethyl cellulose in 150 milliliters of dichloromethane. Then there were added 75 milliliters of dichloromethane and 2.5 milliliters 1,2,3-propanetriol. 10 Grams of polyethylene glycol was molten and dissolved in 75 milliliters of dichloromethane. The latter solution was added to the former and then there were added 2.5 grams of magnesium octadecanoate, 5 grams of polyvinylpyrrolidone and 30 milliliters of concentrated colour suspension (Opaspray K-1-2109) and the whole was homogenated.

The tablet cores were coated with the thus obtained mixture in a coating apparatus.

Example 102 : INJECTABLE SOLUTION

1.8 Grams methyl 4-hydroxybenzoate and 0.2 grams propyl 4-hydroxybenzoate were dissolved in about 0.5 liters of boiling water for injection. After cooling to about 50°C there were added while stirring 4 grams lactic acid, 0.05 propylene glycol and 4 grams of the A.I.. The solution was cooled to room temperature and supplemented with water for injection q.s. ad 1 liter volume, giving a solution of 4 milligrams A.I. per milliliters. The solution was sterilized by filtration (U.S.P. XVII p. 811) and filled in sterile containers.

Example 103 : SUPPOSITORIES


3 Grams A.I. was dissolved in a solution of 3 grams 2,3-dihydroxybutanedioic acid in 25 milliliters polyethylene glycol 400. 12 Grams surfactant and triglycerides q.s. ad 300 grams were molten together. The latter mixture was mixed well with the former solution. The thus obtained mixture was poured onto moulds at a temperature of 37-38°C to form 100 suppositories each containing 30 milligrams of the active ingredient.

The present invention is also related with a method of treating allergic diseases in warm-blooded animals suffering from said allergic diseases by administering an effective anti-allergic amount of a compound of formula (I) or a pharmaceutically acceptable acid addition salt thereof.

Suitable doses administered daily to subjects are varying from 0.1 to 100 mg, more preferably from 1 to 50 mg.

CLAIMS

1. A chemical compound having the formula



- a pharmaceutically acceptable acid addition salt or a possible stereochemically isomeric form thereof, wherein:

- $A^1=A^2-A^3=A^4$ is a bivalent radical having the formula

- $-CH=CH-CH=CH-$ (a-1),
 $-N=CH-CH=CH-$ (a-2),
 $-CH=N-CH=CH-$ (a-3),
 $-CH=CH-N=CH-$ (a-4), or
 $-CH=CH-CH=N-$ (a-5),

- wherein one or two hydrogen atoms in said radicals (a-1) - (a-5) may, each independently from each other, be replaced by halo, lower alkyl, lower alkyloxy, trifluoromethyl or hydroxy;

- R^1 is a member selected from the group consisting of hydrogen, alkyl, cycloalkyl, Ar^1 and lower alkyl substituted with one or two Ar^1 radicals;

- R^2 is a member selected from the group consisting of hydrogen and lower alkyl;

- B is CH_2 , O, S, SO or SO_2 ;

- L is a member selected from the group consisting of a radical of formula

- $L^1-C_{rH_{2r}}-T-C_{sH_{2s}}-$ (b-1); and

- a radical of formula

- $L^1-C_{rH_{2r}}-T^1-N$ (b-2) ;

- wherein one or two hydrogen atoms in the bivalent radical


- $-C_{sH_{2s}}-$ may, each independently from each other, be replaced by

26 halo, hydroxy, mercapto, isothiocyanato, isocyanato, lower alkyloxy,
 27 lower alkylthio, Ar^1 , $\text{Ar}^1\text{O}-$, $\text{Ar}^1\text{S}-$, Ar^1SO_2- , or NR^3R^5 ; and
 28 n is 0 or the integer 1 or 2;
 29 r and s are, independently from each other, 0 or an integer of from
 30 1 to 6 inclusive;

31 T is $-\text{Y}-$ or $-\text{Z}-\overset{\text{X}}{\underset{\parallel}{\text{C}}}-\text{Y}-$;

32 T^1 is $-\text{Z}-\overset{\text{X}}{\underset{\parallel}{\text{C}}}-$ or a direct bond;

33 said Y being O , S , NR^3 or a direct bond;
 34 X being O , S , CH-NO_2 or NR^4 ;
 35 Z being O , S , NR^5 or a direct bond; and
 36 said R^3 being hydrogen, lower alkyl, (Ar^2) lower alkyl, 2-lower
 37 alkyloxy-1,2-dioxoethyl or a radical of formula $-\text{C}(=\text{X})-\text{R}^6$, R^6
 38 being hydrogen, lower alkyl, Ar^2 , Ar^2 -lower alkyl, lower alkyloxy,
 39 Ar^2 -lower alkyloxy, mono- or di(lower alkyl)amino, Ar^2 -amino,
 40 Ar^2 -lower alkylamino or Ar^2 -lower alkyl(lower alkyl)amino;
 41 said R^4 being hydrogen, lower alkyl, cyano, nitro, Ar^2 -sulfonyl,
 42 lower alkylsulfonyl, lower alkylcarbonyl or Ar^2 -carbonyl; and
 43 said R^5 being hydrogen or lower alkyl;
 44 wherein L^1 is a member selected from the group consisting of hydro-
 45 gen; halo; hydroxy; lower alkyloxy; lower alkylthio; cyano; mercapto;
 46 isocyanato; isothiocyanato; Ar^1 ; Ar^1 -carbonyl; Ar^1 -sulfonyl;
 47 lower alkylsulfonyl; cycloalkyl being optionally substituted with up to
 48 two substituents each independently selected from the group consisting
 49 of lower alkyl, cyano and Ar^2 ; [10,11-dihydro-5H-dibenzo[a,d]cyclo-
 50 hepten-5-ylidene]methyl; Het; and furan substituted with substituted
 51 lower alkyl; said substituted lower alkyl being lower alkyl substituted
 52 with a member selected from the group consisting of hydroxy, mercapto,
 53 lower alkyloxy, lower alkylthio, aminolower alkylthio, Ar^2 -oxy and a
 54 radical of formula



- 55 wherein: t is 0 or an integer of from 1 to 6 inclusive; and
 56 R^7 is hydrogen or lower alkyl;
 57 provided that: when in said radical of formula (c) t is
 58 0, then Z or Y is a direct bond; and
- 59 where r is 0, L^1 may also be lower alkenyl, Ar^1 -lower alkenyl or
 60 lower alkyl substituted with two lower alkyloxy radicals; and
- 61 where r is 0 and T is NR^3 , or T is $-N(R^5)-C(=X)-Y$ or T^1 is $-N(R^5)-C(=X)-$,
 62 L^1 may also be amino, lower alkylamino or Ar^1 -amino; and
- 63 where r is 0, and T is $-N(R^5)-C(=X)-Y$ or T^1 is $-N(R^5)-C(=X)-$,
 64 L^1 may also be nitro;
- 65 said Het being an optionally substituted five- or six-membered hetero-
 66 cyclic ring, being optionally condensed with an optionally substituted
 67 five- or six-membered carbocyclic or heterocyclic ring;
- 68 provided that:
- 69 i) when L is a radical of formula (b-1) wherein L^1 is hydrogen and
 70 wherein T is $-Z-C(=X)-Y-$ wherein Y is other than a direct bond
 71 and Z and X are each independently O or S, then r is not 0; or
 72 when L is a radical of formula (b-2) wherein L^1 is hydrogen and
 73 wherein T^1 is $-Z-C(=X)-$ wherein Z and X are each independently
 74 O or S, then r is not 0;
- 75 ii) when L is a radical of formula (b-1) wherein L^1 is halo,
 76 hydroxy, lower alkyloxy, mercapto, lower alkylthio, isocyanato,
 77 isothiocyanato or Het connected to C_rH_{2r} on a nitrogen atom,
 78 and wherein r is 0, then T is a direct bond or a radical
 79 $-C(=X)-Y-$; or when L is a radical of formula (b-2) wherein L^1
 80 is halo, hydroxy, lower alkyloxy, mercapto, lower alkylthio,
 81 isocyanato, isothiocyanato or Het connected to C_rH_{2r} on a
 82 nitrogen atom, and wherein r is 0, then T^1 is a radical $-C(=X)-$;
- 83 iii) when L is a radical of formula (b-1) wherein T is Y, said Y being
 84 other than a direct bond, or wherein T is $-Z-C(=X)-Y-$, wherein Y

85 is other than a direct bond, then s is not 0;


86 wherein Ar^1 is a member selected from the group consisting of
 87 phenyl, substituted phenyl, naphthalenyl, thienyl, halothienyl, lower
 88 alkylthienyl, pyridinyl, mono- and di(lower alkyloxy)pyridinyl,
 89 pyrrolyl, lower alkylpyrrolyl, furanyl, furanyl substituted with lower
 90 alkyl, pyrazinyl, thiazolyl, imidazolyl, lower alkylimidazolyl;
 91 said substituted phenyl, being phenyl substituted with up to 3
 92 substituents each independently selected from the group consisting of
 93 halo, hydroxy, nitro, cyano, trifluoromethyl, lower alkyl, lower
 94 alkyloxy, lower alkylthio, mercapto, amino, mono- and di(lower
 95 alkyl)amino, lower alkylsulfonyl, lower alkylsulfonylower alkyl,
 96 phenyllower alkylsulfonyl, phenylsulfonylower alkyl, a radical of
 97 formula $R^8-C_{p-2p}H_{2p}-Y-$, a radical of formula $R^9-Z-C(=X)-Y-$, and a
 98 radical of formula $R^{10}SO_2Y-$; wherein p is an integer of from 1 to
 99 6 inclusive and R^8 is a member selected from the group consisting of
 100 amino, cyano, phenyl aminocarbonyl, mono- and di(lower alkyl)amino-
 101 carbonyl, lower alkyloxycarbonyl, phenyllower alkyloxycarbonyl,
 102 4-morpholinylcarbonyl, 1-piperidinylcarbonyl, 1-pyrrolidinylcarbonyl,
 103 and lower alkenyl; wherein R^9 is member selected from the group
 104 consisting of hydrogen, lower alkyl and Ar^2 ; provided that, when
 105 R^9 is hydrogen and Y is other than a direct bond, then Z is not O or
 106 S; and wherein R^{10} is lower alkyl or Ar^2 ;

107 wherein Ar^2 is a member selected from the group consisting of
 108 phenyl, substituted phenyl, thienyl and furanyl, said substituted
 109 phenyl being phenyl optionally substituted with up to three
 110 substituents each independently selected from the group consisting of
 111 halo, hydroxy, nitro, cyano, trifluoromethyl, lower alkyl, lower
 112 alkyloxy, lower alkylthio, mercapto, amino, mono- and di(lower
 113 alkyl)amino, carboxyl, lower alkyloxycarbonyl and (lower alkyl)-CO.


1 2. A compound according to claim 1, wherein r is 0 and L^1 is
 2 hydrogen, hydroxy, lower alkyloxy, lower alkylthio, mercapto, Het,
 3 Ar^1 , cyanato, isothiocyanato, or isocyanato.

1 3. A compound according to claim 2, wherein R^1 is lower alkyl
 2 substituted with one Ar^1 radical.


- 1 4. A compound according to claim 3, wh r in L is a radical of
2 formula (b-1).
- 1 5. A chemical compound according to any of claims 1 to 4 for use
2 as a medicine.
- 1 6. A chemical compound according to any of claims 1 to 4 for use
2 as an anti-allergic medicine.
- 1 7. A pharmaceutical composition comprising an inert carrier and a
2 pharmaceutically acceptable amount of a compound according to any of
3 claims 1 to 4.
- 1 8. A pharmaceutical composition according to claim 7 for use as an
2 anti-allergic medicine.
- 1 9. A process for preparing a pharmaceutical composition,
2 characterized in that a therapeutically effective amount of a
3 compound as claimed in any of claims 1 to 4 is intimatately mixed
4 with a suitable pharmaceutical carrier.
- 1 10. A process for preparing a compound according to any of claims
2 1 to 4 characterized by
3 I reacting a piperidine of formula




- 4 wherein X^1 is O, S or NH and W is a reactive leaving group, with
5 a diamine of formula



- 6 in a reaction-inert solvent;
- 7 II reacting a piperidine of formula



9




(V), in a reaction-inert

10 solvent, wherein :


- 11 a) E^1 is a radical of formula $-B-M$ wherein M is hydrogen or an
 12 alkalimetal or earth alkalimetal and E^2 is a radical of
 13 formula $-W$; or
 14 b) E^1 is a radical of formula $-W$ and E^2 is a radical of
 15 formula $M-B$; or
 16 c) E^1 is a radical of formula $-CH_2-W$ and E^2 is a radical of
 17 formula $-M$, thus preparing a compound of formula

18



(I-a); or


- 19 d) E^1 is a radical of formula $-M$ and E^2 is a radical of
 20 formula $-CH_2W$, thus preparing a compound of formula (I-a);
 21 III. reducing an intermediate of formula




, (VI)

- 22 in a reaction-inert solvent; and, if desired, converting the
 23 compounds of formula (I) into each other by
 24 a) alkylating a compound of formula Q^2-D (I-c) with a reagent of
 25 formula L^1-Q^1 (VII) in a suitable solvent, thus preparing a
 26 compound of formula L^2-D (I-b), wherein L^2 has the
 27 previously defined meaning of L , provided that it is other than
 28 hydrogen, and wherein
 29 1) Q^1 is $-W$ and Q^2 is hydrogen; or

- 30 ii) Q^1 is $-C_{rH_{2r}}-W^1$ and Q^2 is a radical of formula
 31 $HT^2-C_{sH_{2s}}-$, wherein W^1 is a reactive leaving group
 32 and T^2 is O, S, NR^3 or $-Z^1-C(=X)-Y-$, said Z^1 being
 33 O, S or NR^5 , thus preparing a compound of formula
 34 $L^1-C_{rH_{2r}}-T^2-C_{sH_{2s}}-D$ (I-b-1-a); or
 35 iii) Q^1 is $-C_{rH_{2r}}-W^1$ and Q^2 is a radical of formula



37 or $Z^1-C(=X)-$, thus preparing a compound of formula



- 39 iv) Q^1 is a radical of formula $-C_{rH_{2r}}-T^4H$ and Q^2 is
 40 $W-C_{sH_{2s}}-$, wherein T^4 is O, S, NR^3 or $-Z-C(=X)-Y^1-$,
 41 said Y^1 being O, S or NR^3 , thus preparing a compound of
 42 formula $L^1-C_{rH_{2r}}-T^4-C_{sH_{2s}}-D$ (I-b-2);

43 b) reductively N-alkylating a compound of formula H-D, (I-c-1),
 44 with a carbonyl-compound of formula $L^{2-a}=C=O$ (VIII), said
 45 $L^{2-a}=C=O$ being a compound of formula L^2-H wherein a $-CH_2-$
 46 radical is oxidated to a carbonyl radical, in a reaction-inert
 47 solvent, thus preparing a compound of formula L^2-D (I-b);

48 c) reductively N-alkylating a compound of formula



50 with a carbonyl-compound of formula $L^1-(C_{rH_{2r-1}})=O$ (IX),

51 said $L^1-(C_{rH_{2r-1}})=O$ being a compound of formula

52 $L^1-C_{rH_{2r}}-H$ wherein a $-CH_2-$ radical is oxidated to a

53 carbonyl radical, in a reaction inert solvent, thus preparing a

54 compound of formula $L^1-C_{rH_{2r}}-N(R^3)-C_{sH_{2s}}-D$ (I-b-3);

55 d) reductively N-alkylating an intermediate of formula


56 $L^1-C_{rH_{2r}}-N(R^3)H$ (X) with a compound of formula $O=(C_{sH_{2s-1}})-D$

57 (I-e), said $O=(C_{sH_{2s-1}})-$ being a radical of formula $H-C_{sH_{2s}}-$


58 wherein a $-CH_2-$ radical is oxidated to a carbonyl radical, in a

59 reaction-inert solvent, thus preparing a compound of formula (I-b-3);


- 60 e) reacting a reagent of formula $L^1-C_rH_{2r}-Z^1H$ (XI) with a
 61 compound of formula $X^2=C=N-C_sH_{2s}-D$ (I-f), wherein X^2 is
 62 O or S, in a reaction-inert solvent, thus preparing a compound
 63 of formula $L^1-C_rH_{2r}-Z^1-C(=X^2)-NH-C_sH_{2s}-D$ (I-b-4);
 64 f) reacting a reagent of formula $L^1-C_rH_{2r}-N=C=X^2$ (XII) with
 65 a compound of formula $HY^1-C_sH_{2s}-D$ (I-c-4), respectively
 66 with a compound of formula $H-D$ (I-c-1) or with a compound of
 67 formula



- 68 (I-c-5), in a reaction-inert solvent, thus
 69 preparing a compound of formula




- 72 respectively of formula




- 74 $L^1-C_rH_{2r}-NH-C(=X^2)-N$ (I-b-5-c);

- 75 g) reacting a reagent of formula $L^1-C_rH_{2r}-C(=X^2)-OH$ (XIII)
 76 with (I-c-4), respectively with (I-c-1) or (I-c-5), in a
 77 reaction-inert solvent, if desired, after converting the OH
 78 group in (XIII) in a reactive leaving group or by reacting
 79 (XIII) with (I-c-4), respectively (I-c-1) or (I-c-5), in the
 80 presence of a reagent capable of forming esters or amides, thus
 81 preparing a compound of formula




- 83 respectively of formula





- 85 $L^1-C_rH_{2r}-C(=X^2)-N$ (I-b-6-c);

- 86
 87 h) reacting (XI) with (I-c-4), respectively (I-c-1) or (I-c-5) in
 88 the presence of a $C=X$ generating agent in a reaction-inert
 89 solvent, thus preparing a compound of formula




- 91 respectively of formula







- i) reacting an alkene of formula $L^1-C_{rH_{2r}}-T$ -lower alkenediyl-H of formula (XIV) with (I-c-1) in a reaction-inert solvent, thus preparing a compound of formula



- j) reacting a reagent of formula




in a reaction-inert solvent, thus preparing a compound of formula




wherein s' is an integer of from 2 to 6 inclusive;

- k) cyclizing an imidamide of formula




in a reaction-inert solvent in the presence of an acid, thus preparing a compound of formula




and R^{23} are, each independently, optional substituents of the imidazole ring;


- l) condensing a ketone of formula $R^{24}-CH(W)-C(=O)-R^{25}$, (XVII), with a thioamide of formula $H_2N-C(=S)-K-D$, (XVIII), in a reaction-inert solvent, thus preparing a compound of formula




- 115 R^{25} are, each independently, optional substituents of the
 116 thiazole ring, or where in the compound of formula (I-i-2) said
 117 thiazolyl ring is condensed with a five- or six-membered hetero-
 118 or carbocyclic ring, R^{24} and R^{25} taken together may form a
 119 radical of formula G^3 ;
 120 m) condensing a thioamide of formula $R^{26}-C(=S)NH_2$, (XIX), with
 121 a ketone of formula $W-CHR^{27}-(C=O)-K-D$, (XX), in a reaction-
 122 inert solvent, thus preparing a compound of formula




- 124 R^{27} are, each independently, optional substituents of the
 125 thiazolyl ring;
 126 n) reacting an amide or thioamide of formula



- 128 generating agent, in a reaction-inert solvent, thus preparing a
 129 compound of formula




- 130 o) cyclizing a urea or thiourea of formula



- 131 which in situ may be generated by reacting a reagent

132




(XXIII), with an amine

133

H₂N-K-D, (XXIV), in a reaction-inert solvent, thus preparing a compound of formula

134


135



(I-i-4-a);

136

p) condensing an aniline of formula



(XXV),

137

with an acid of formula R¹²COOH (XXVI), or a reactive


138

derivative thereof, in a reaction-inert solvent, thus preparing

139

a compound of formula

140




(I-i-5);

141

q) condensing an aniline of formula

142



(XXVII), with an amide

143


of formula R¹²-C(=O)-NH-K-D (XXVIII) in a reaction-inert

144

solvent, thus preparing a compound of formula (I-i-5);


145

r) condensing an aniline of formula




(XXIX),


146 with an acetylene of formula $\text{CH}\equiv\text{C}-\text{R}^{14}$ (XXX), in a reaction-
 147 inert solvent, thus preparing a compound of formula




148 s) condensing (XXIX) with a ketone of formula $\text{R}^{14}-\text{C}(=\text{O})-\text{R}^{15}$
 149 (XXXI), in a reaction-inert solvent, thus preparing a compound
 150 of formula




151 t) condensing a reagent of formula




152 with a ketone of formula $\text{W}-\text{CH}(\text{R}^{16})-\text{C}(=\text{O})-\text{K}-\text{D}$ (XXXIII), in a
 153 reaction-inert solvent, thus preparing a compound of formula




154 u) condensing an amine of formula




155 with CS_2 , in a reaction-inert solvent, thus preparing a
 156 compound of formula




- 157 v) reacting a reagent of formula $R^{19}-C(=NH)-W$ (XXXV) with an
 158 amine of formula




- 159 in a reaction-inert solvent, thus preparing a compound of formula




- 160 w) cyclodesulfurizing a thioamide of formula




- 161 with an appropriate alkyl halide, metal oxide or metal salt in a
 162 reaction-inert solvent, thus preparing a compound of formula



- 163 x) condensing an amine of formula





- 164 with a $>C=X^2$ generating agent, in a reaction-inert solvent,
 165 thus preparing a compound of formula




(I-1-12);

166 wherein K is a bivalent radical of formula



169 wherein D represents a radical of formula



170 $R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}$ and R^{20} are, each
 171 independently optional substituents of the previously described
 172 bicyclic radicals and G^1, G^2, G^3, G^4 and G^5 are, each independently,
 173 optionally substituted bivalent radicals, selected so that they form,
 174 combined with the five- or six-membered heterocycle to which they are
 175 attached, a bicyclic Het-system; or optionally converting the
 176 compounds of formula (I) into each other following art-known
 177 grouptransformation procedures, and, if desired, converting the
 178 compounds of formula (I) into a therapeutically active non-toxic
 179 acid-addition salt form by treatment with an appropriate acid or,
 180 conversely, converting the acid-addition salt into the free base form
 181 with alkali; and/or preparing stereochemically isomeric forms thereof.



DOCUMENTS CONSIDERED TO BE RELEVANT			EP 84201851.7
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
D,A	<p><u>US - A - 4 219 559</u> (JANSSENS)</p> <p>* Column 1, formula I; column 3, lines 24-39; column 11, lines 62-65; column 12, lines 33-35 *</p> <p>--</p>	1,5-10	<p>C 07 D 471/04</p> <p>C 07 D 401/06</p> <p>C 07 D 401/12</p> <p>C 07 D 401/14</p> <p>C 07 D 405/14</p> <p>C 07 D 409/14</p> <p>C 07 D 417/14</p> <p>C 07 D 513/04</p>
A	<p><u>GB - A - 1 260 857</u> (HEXACHIMIE)</p> <p>* Page 1, lines 1-44; claim 7 *</p> <p>--</p>	1,5-8, 10	<p>C 07 D 519/00</p> <p>C 07 D 487/04</p> <p>C 07 D 473/00</p> <p>C 07 D 473/34</p> <p>A 61 K 31/445</p>
A	<p><u>US - A - 2 857 391</u> (ENGELHARDT)</p> <p>* Column 1, lines 23-30, 58-60 *</p> <p>--</p>	1,5	
A	<p>CHEMICAL ABSTRACTS, vol. 96, no. 9, March 1, 1982, Columbus, Ohio, USA</p> <p>P.B. BERNTSSON "Heterocyclic compounds useful for inducing stomach acid secretion" page 613, column 1, abstract no. 68 986p</p> <p>& Swed.SE 416 649, 26 Jan. 1981</p> <p>--</p>	1,5	<p>TECHNICAL FIELDS SEARCHED (Int. Cl. 4)</p> <p>C 07 D 471/00</p> <p>C 07 D 401/00</p> <p>C 07 D 405/00</p> <p>C 07 D 409/00</p> <p>C 07 D 417/00</p> <p>C 07 D 513/00</p> <p>C 07 D 519/00</p> <p>C 07 D 487/00</p> <p>C 07 D 473/00</p>
A	<p><u>AT - B - 332 426</u> (BOEHRINGER MANNHEIM)</p> <p>* Page 1, formula I, lines 8,9 *</p> <p>--</p>	1,5-8	
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
VIENNA		10-04-1985	ONDER
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technical background</p> <p>O : non-written disclosure</p> <p>P : intermediate document</p> <p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

EPO Form 1503 03/82



DOCUMENTS CONSIDERED TO BE RELEVANT			EP 84201851.7
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	<p>EP - B1 - 0 045 251 (RHONE-POULENC)</p> <p>* Page 2, formula I; page 3, lines 43-45 *</p> <p>-----</p>	1, 5-8	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 10-04-1985	Examiner ONDER
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO Form 1503 (03/82)

